



Eriophyoid mite fauna (Acari: Eriophyoidea) of Maragheh county in Iran

Nesa Yousefnezhad¹ | Maghsoud Pazhouhandeh² | Parisa Lotfollahi^{1*}

1. Department of Plant Protection, Faculty of Agriculture, Azarbaijan Shahid Madani University, Tabriz, Iran; E-mails: nesayousefnezhad@gmail.com, prslotfollahi@yahoo.com

2. Department of Biotechnology, Faculty of Agriculture, Azarbaijan Shahid Madani University, Tabriz, Iran; E-mail: pazhouhandeh@gmail.com

* Correspondence

✉ prslotfollahi@yahoo.com

Received:
09 August, 2025

Accepted:
29 August, 2025

Published:
15 October, 2025

Subject Editor:
A. Saboori



CITE: Yousefnezhad, N., Pazhouhandeh, M. & Lotfollahi, P. (2025) Eriophyoid mite fauna (Acari: Eriophyoidea) of Maragheh county in Iran. *Persian Journal of Acarology*, 14(4): 140412.

<https://doi.org/10.22073/pja.v14i4.87544>

Mites of the superfamily Eriophyoidea Nalepa (Acari: Acariformes) are obligate plant parasites with high host specialization and are the second most important group of mites causing damage to agricultural crops after the superfamily Tetranychidae Lindqvist (Lindquist *et al.* 1996; Krantz and Walter 2009). Mites of this superfamily, with a global distribution, are known as bud mites, blister mites, erinose mites, rust mites and four-legged mites, and are currently included in three families: Eriophyidae Nalepa, Diptilomiopidae Keifer and Phytoseptidae Murray. These mites feed on all aerial parts of annual, biennial and perennial plants and some ferns except the roots. Members of this superfamily account for more than 60% of herbivorous mites (Li *et al.* 2023). In addition to the direct damage caused by eriophyoid mites to agricultural crops, their ability to transmit plant viruses and their potential as biological control agents for some weeds are of great importance (Oldfield 2005). Therefore, because of their high importance, the eriophyoid mite fauna was surveyed in 15 locations of Maragheh county, East Azerbaijan province, Northwest of Iran, in summers 2023 and 2024 (Table 1; see supplement for Tables). Eriophyoid mites were recovered from the plant material by means of a modified washing method developed by Lotfollahi and Masoudi-Rad (2024). The collected specimens were slide mounted according to Lotfollahi and Masoudi-Rad (2024). In order to make the species identification possible, all measurements were taken using an Olympus BX53 microscope, through a phase contrast 100× oil immersion objective at 1,000 magnification, according to Amrine and Manson (1996) as modified by de Lillo *et al.* (2010), and counting of dorsal, ventral and coxigenital semiannuli follows Lotfollahi and Masoudi-Rad (2024). The genera classification follows Amrine *et al.* (2003) and comparisons were also made with new genera described since that publication. The examined materials are deposited at the Acarology Laboratory, Department of Plant Protection, Faculty of Agriculture, Azarbaijan Shahid Madani University, Tabriz (Iran). Host plant names and their synonymies are in accordance with "*The World Flora Online*" (2025).

In total specimens belonging to 12 species, two families, three subfamilies, four tribes, and 9 genera were collected and identified, and the information about them are provided herein. In this study, the highest number of mite specimens belonged to *Calepitrimerus baileyi* Keifer, 1938 and the lowest number belonged to the *cynodonis* Abou-Awad & Nasr, 1983.

Family Eriophyidae Nalepa

Subfamily Eriophyinae Nalepa

Tribe Acerini Amrine & Stasney

Genus *Aceria* Keifer



***Aceria lactulignae* Honarmand & Lotfollahi, 2024**

Locality and host plant – Collected by N. Yousefnezhad during summer 2024; on plants and localities listed in Table 2.

Previous provincial records for Iran – East Azerbaijan (Honarmand and Lotfollahi 2024).

Remarks – It should be supposed that this mite species can be subjected to wind dispersal and can be landed on non-elective host plant species, considering the collection of solely few females. This is a new record to the fauna of Maragheh County.

***Aceria xeranthemis* Lotfollahi, Haddad, Khanjani, Moghaddam & de Lillo, 2013**

Locality and host plant – Collected by N. Yousefnezhad during summers 2023 and 2024; on plants and localities listed in Table 3.

Previous provincial records for Iran – Ardabil, East Azerbaijan (Lotfollahi *et al.* 2013; Hemmatzadeh-khorshidabadi *et al.* 2023)

Remarks – No apparent symptom was observed on the elective host plants.

Subfamily Phyllocoptinae Nalepa**Tribe Tegenotini Bagdasarian****Genus *Shevtchenkella* Bagdasarian*****Shevtchenkella denticulata* Lotfollahi, de Lillo & Haddad, 2014a**

Locality and host plant – Collected by N. Yousefnezhad during summer 2024; on plants and localities listed in Table 5.

Previous provincial records for Iran – East Azerbaijan, West Azerbaijan (Lotfollahi *et al.* 2014a, 2023).

Remarks – No apparent symptom was observed on the host plant. This is a new record to the fauna of Maragheh County.

Tribe Phyllocoptini Nalepa**Genus *Phyllocoptes* Nalepa*****Phyllocoptes abaenus* Keifer, 1940**

Locality and host plant – Collected by N. Yousefnezhad during summer 2023; on plants and localities listed in Table 6.

Previous provincial records for Iran – Ardabil, East Azerbaijan, Hamedan, Mazandaran, West Azerbaijan (Lotfollahi *et al.* 2014b; Gol *et al.* 2018; Mehri-Heyran *et al.* 2020; Jafari and Khanjani 2021; Shalchi-zonouz *et al.* 2022; Hemmatzadeh-khorshidabadi *et al.* 2023; Lotfollahi *et al.* 2023; Ranjbar-Varandi *et al.* 2023).

Remarks – It should be supposed that this mite species can be subjected to wind dispersal and can be landed on non-elective host plant species, considering the collection of solely few females. No apparent symptom was observed on the elective and non-elective host plants. This is a new record to the fauna of Maragheh County.

Genus *Calepitrimerus* Keifer***Calepitrimerus achilleae* Roivainen, 1950**

Locality and host plant – Collected by N. Yousefnezhad during summer 2024; on plants and localities listed in Table 7.

Previous provincial records for Iran – No previous record.

Remarks – No apparent symptom was observed on the elective host plant. This is a new record to the fauna of Iran.

Calepitrimerus baileyi Keifer, 1938

Locality and host plant – Collected by N. Yousefnezhad during summers 2023 and 2024; on plants and localities listed in Table 8.

Previous provincial records for Iran – Ardabil, East Azerbaijan, Golestan, Lorestan, West Azerbaijan (Lotfollahi *et al.* 2014b; Gol *et al.* 2018; Mehri-Heyran *et al.* 2020; Bahirai *et al.* 2021; Hemmatzadeh-khorshidabadi *et al.* 2023; Lotfollahi *et al.* 2023).

Remarks – It should be supposed that this mite species can be subjected to wind dispersal and can be landed on non-elective host plant species, considering the collection of solely few females. No apparent symptom was observed on the elective and non-elective host plants. This is a new record to the fauna of Maragheh County.

Tribe Anthoceptini Amrine & Stasny

Genus *Aculus* Keifer

Aculus fockeui (Nalepa & Trouessart, 1891)

Locality and host plant – Collected by N. Yousefnezhad during summer 2023; on plants and localities listed in Table 9.

Previous provincial records for Iran – Ardabil, East Azerbaijan, Hamedan, Lorestan, West Azerbaijan (Lotfollahi *et al.* 2014b; Mehri-Heyran *et al.* 2020; Bahirai *et al.* 2021; Jafari and Khanjani 2021; Shalchi-Zonouz *et al.* 2022; Hemmatzadeh-khorshidabadi *et al.* 2023; Lotfollahi *et al.* 2023).

Remarks – It should be supposed that this mite species can be subjected to wind dispersal and can be landed on non-selective host plant species, considering the collection of solely few females.

Aculus comatus (Nalepa, 1892)

Locality and host plant – Collected by N. Yousefnezhad during summer 2023; on plants and localities listed in Table 10.

Previous provincial records for Iran – East Azerbaijan (Bagheri *et al.* 2024).

Remarks – No apparent symptom was observed on the elective host plant. This is a new record to the fauna of Maragheh County.

Genus *Aculops* Keifer

Aculops pachyphora Honarmand, Sadeghi & de Lillo, 2020

Locality and host plant – Collected by N. Yousefnezhad during summer 2023; on plants and localities listed in Table 11.

Previous provincial records for Iran – South Khorasan, West Azerbaijan (Honarmand *et al.* 2020; Lotfollahi *et al.* 2023).

Remarks – No apparent symptom was observed on the elective host plant. This is a new record to the fauna of East Azerbaijan province.

Genus *Anthoceptes* Nalepa

Anthoceptes salicis (Nalepa, 1894)

Locality and host plant – Collected by N. Yousefnezhad during summer 2023; on plants and localities listed in Table 12.

Previous provincial records for Iran – East Azerbaijan, Lorestan, Mazandaran, Zanjan (Tarasi and Taghaddosi 2005, Lotfollahi *et al.* 2014c, Bahirai *et al.* 2021; Ranjbar-Varandi *et al.* 2023).

Remarks – It should be supposed that this mite species can be subjected to wind dispersal and can be landed on non-elective host plant species. This is a new record to the fauna of Maragheh County.

Genus *Abacarus* Keifer

***Abacarus cynodonis* Abou-Awad et Nasr, 1983**

Locality and host plant – Collected by N. Yousefnezhad during summer 2023; on plants and localities listed in Table 13.

Previous provincial records for Iran – West Azerbaijan (Lotfollahi *et al.* 2023).

Remarks – It should be supposed that this mite species can be subjected to wind dispersal and can be landed on non-elective host plant species, considering the collection of solely one female. This is a new record to the fauna of East Azerbaijan province.

Family Diptilomiopidae Keifer

Subfamily Rhyncaphyoptinae Roivainen

Genus *Rhinophytoptus* Liro

***Rhinophytoptus nemalobos* Lotfollahi & de Lillo, 2014**

Locality and host plant – Collected by N. Yousefnezhad during summer 2023; on plants and localities listed in Table 15.

Previous provincial records for Iran – East Azerbaijan, Hamedan, Lorestan, Mazandaran, West Azerbaijan (Lotfollahi *et al.* 2014b; Mehri-Heyran *et al.* 2020; Jafari and Khanjani 2021; Shalchi-Zonouz *et al.* 2022; Ranjbar-Varandi *et al.* 2023).

Remarks – It should be supposed that this mite species can be subjected to wind dispersal and can be landed on non-elective host plant species, considering the collection of solely few females and males. No apparent symptom was observed on the elective and non-elective host plants. This is a new record to the fauna of Maragheh County.

ACKNOWLEDGEMENTS

This research was partially supported by Azarbaijan Shahid Madani University and Iran National Science Foundation (4001458).

Author contributions: Methodology: P.L., M.P.; Sampling, collection and preparation of specimens: N.Y.; Analyses and data curation: P.L., M.P.; Illustration and photography: N.Y.; writing – original draft preparation: N.Y.; writing – revisions and editing: P.L., M.P.; supervision: P.L.; funding acquisition: P.L. All authors have read and agreed to the published version of the manuscript.

Funding: The research was funded by the Azarbaijan Shahid Madani University and Iran National Science Foundation (Grant No. 4001458).

Data availability: Data are available upon request from the authors.

Ethics approval: This study only included arthropod material, and all required ethical guidelines for the treatment and use of animals were strictly adhered to in accordance with international, national, and institutional regulations. No human participants were involved in any studies conducted by the authors for this article.

Conflict of interest: The authors declare no conflict of interest.

REFERENCES

Abou-Awad, B.A. & Nasr, A.K. (1983) Occurrence of the eriophyid mites as new pests of Bermuda grass, *Cynodon*

dactylon (L.) Pers., in Egypt. *International Journal of Acarology*, 9(4): 183–187.

- Amrine, J.W. Jr. & Manson, D.C.M. (1996) Preparation, mounting and descriptive study of eriophyoid mites. In: Lindquist, E.E., Sabelis, M.W. & Bruin, J. (Eds.), *Eriophyoid mites – their biology, natural enemies and control. World Crop Pests, Vol. 6*. Elsevier Science, pp. 383–396. [https://doi.org/10.1016/S1572-4379\(96\)80023-6](https://doi.org/10.1016/S1572-4379(96)80023-6)
- Amrine, J.W. Jr., Stasny, T.A.H. & Flechtmann, C.H.W. (2003) *Revised keys to world genera of Eriophyoidea (Acari: Prostigmata)*. West Bloomfield, Michigan, USA, Indira Publishing House, 244 pp.
- Bagheri, M., Maghsoodi, S., Jahandideh-Dogijan, J. & Lotfollahi, P. (2024) Eriophyoid mites of East Bostanabad in East Azerbaijan province. *Proceeding of 25th Iranian Plant Protection Congress, 7–10 September, IROST, Tebran*, pp. 665–666.
- Bahirai, F., Jafari, S., Lotfollahi, P. & Shakarami, J. (2021) Eriophyoidea (Acari: Trombidiformes) of the Lorestan Province and first record of *Aceria querci* (Garnam, 1883) outside of the USA. *Persian Journal of Acarology*, 10(1): 111–119. <https://doi.org/10.22073/pja.v10i1.63306>
- de Lillo, E., Craemer, C., Amrine, J.W. Jr. & Nuzzaci, G. (2010) Recommended procedures and techniques for morphological studies of Eriophyoidea (Acari: Prostigmata). In: Ueckermann, E.A. (Ed.) *Eriophyoid mites: progress and prognoses*. Springer, Dordrecht, pp. 283–307. https://doi.org/10.1007/978-90-481-9562-6_15
- Gol, A., Sadeghi-Nameghi, H. & de Lillo, E. (2018) Investigation of eriophyid mites (Acari: Trombidiformes: Eriophyoidea) of trees and shrubs in Golestan province. *Proceedings of the 23rd Iranian Plant Protection Congress, 27-30 Aug., Gorgan University of Agricultural Sciences and Natural Resources*, p. 1636.
- Hemmatzadeh-khorshidabadi, H., Lotfollahi, P., Mehrvar, A., Shiri, J. & de Lillo, E. (2023) Eriophyoid (Acari: Eriophyoidea) mite fauna of Hesar village in Meshginshahr region with description of a new species. *Persian Journal of Acarology*, 12(3): 403–415. <https://doi.org/10.22073/pja.v12i3.81117>
- Honarmand, A. & Lotfollahi, P. (2024) *Aceria lactulignae* sp. nov. (Acari Eriophyidae) from Northwest Iran. *Redia*, 107: 43–46. <https://doi.org/10.19263/REDIA-107.24.06>
- Honarmand, A., Sadeghi-Namaghi, H. & de Lillo, E. (2020) Three new species of eriophyid mites (Trombidiformes: Eriophyoidea) associated with Leguminosae species from semi-arid and arid environment in East Iran. *International Journal of Acarology*, 46(4): 201–207. <https://doi.org/10.1080/01647954.2020.1752797>
- Jafari, S. & Khanjani, M. (2021) Five new eriophyoid mite species associated with plum orchards, *Prunus domestica* L. and five new records of the superfamily Eriophyoidea (Acari: Trombidiformes) from Hamedan Province, Iran. *Systematic and Applied Acarology*, 26(8): 1543–1559. <https://doi.org/10.11158/saa.26.8.11>
- Keifer, H.H. (1938) Eriophyid Studies II. *Bulletin of the California Department of Agriculture*, 27: 301–323.
- Keifer, H.H. (1940) Eriophyid Studies VIII. *Bulletin of the California Department of Agriculture*, 29: 21–46.
- Krantz, G.W. & Walter, D.E. (Eds.) (2009) *A Manual of Acarology*. Third Edition. Texas Tech University Press, USA, 816 pp.
- Li, N., Sun, J.-T., Yin, Y., Hong, X.-Y. & Xue, X.-F. (2023) Global patterns and drivers of herbivorous eriophyoid mite species diversity. *Journal of Biogeography*, 50(2): 330–340. <https://doi.org/10.1111/jbi.14782>
- Lindquist, E.E., Sabelis, M.W., Bruin, J. (Eds.) (1996) *Eriophyoid Mites: Their Biology, Natural Enemies and Control*. World Crop Pests, Vol. 6, Elsevier Science Publishers, Amsterdam, Netherlands, 760 pp.
- Lotfollahi, P. & Masoudi-Rad, S. (2024) One new *Aceria* species (Acari: Eriophyoidea) from Jolfa, Iran. *Persian Journal of Acarology*, 13(2): 169–176. <https://doi.org/10.22073/pja.v13i2.84596>
- Lotfollahi, P., Aghazadeh, K. & Azimi, S. (2023) Mite fauna of subfamily Phyllocoptinae (Acari Eriophyidae) in Maku county with three new records for Asia and three new records for Iran. *Redia*, 106: 107–115. <https://doi.org/10.19263/REDIA-106.23.12>
- Lotfollahi, P., de Lillo, E. & Haddad Irani-Nejad, K. (2014a) Three new species from the subfamily Phyllocoptinae (Acari, Trombidiformes, Eriophyidae) in Iran. *ZooKeys*, 426: 17–27.
- Lotfollahi, P., Haddad Irani-Nejad, K. & de Lillo, E. (2014b) Eriophyoid mites (Acari: Trombidiformes: Eriophyoidea) of Rosales trees in Iran: two new species and three new records. *Zootaxa*, 3861(1): 76–85. <https://doi.org/10.11646/zootaxa.3861.1.4>
- Lotfollahi, P., Haddad Irani-Nejad, K. & de Lillo, E. (2014c) Eight new records for the eriophyid (Trombidiformes Eriophyoidea Eriophyidae) mite fauna of Iran. *Redia*, 97: 51–61.
- Lotfollahi, P., Haddad Irani-Nejad, K., Khanjani, M., Maghadam, M. & de Lillo, E. (2013) Eriophyoid mites (Acari: Prostigmata: Eriophyidae) associated with Compositae in Iran. *Zootaxa*, 3664(3): 349–360. <https://doi.org/>

10.11646/zootaxa.3664.3.5

- Mehri-Heyran, H., Lotfollahi, P., de Lillo, E. & Azimi, S. (2020) Eriophyoid (Trombidiformes: Eriophyoidea) mite fauna of Miandoab region in Iran with redescription of *Aceria kiefferi* (Nalepa). *Persian Journal of Acarology*, 9(2): 161–171. <https://doi.org/10.22073/pja.v9i2.59382>
- Nalepa, A. (1891) Diagnoses d'acariens nouveaux Le Naturaliste. *Revue Illustrée des Sciences Naturelles Sér. 2*, 13(93): 25–26 (Paris) [=Trouessart 1891].
- Nalepa, A. (1892) Neue Gallmilben. 5. *Fort. Anzeiger der Kaiserlichen Akademie Wissenschaften. Mathematisch-naturwissenschaftliche Klasse*, Wien., 29(19): 190–192.
- Nalepa, A. (1894) Beiträge zur Kenntnis der Phyllocoptiden. *Nova Acta Leopoldino-Carolinae Deutschen Akademie der Naturforscher*, 61(4): 289–324 + 6 pls.
- Oldfield, G.N. (2005) Biology of gall-inducing Acari. In: Raman, A., Schaefer, C.W. & Withers, T.M. (Eds.), *Biology, Ecology and Evolution of Gall-Inducing Arthropods*. Science Publishers, Inc., Enfield (NH), USA, pp. 35–57.
- Ranjbar-Varandi, F., Haddad Irani-Nejad, K. & Lotfollahi, P. (2023) Eriophyoid mite fauna (Acari: Eriophyoidea) of Sari county in Iran with seven new records for Mazandaran province. *Persian Journal of Acarology*, 12(4): 581–586. <https://doi.org/10.22073/pja.v12i4.82955>
- Roivainen, H. (1950) Eriophyid news from Sweden. *Acta Entomologica Fennica*, 7: 1–51.
- Shalchi-Zonouz, M., Ataie-Esfahlan, S., Haghverdizadeh, S., Amjadi-Shadbad, N. & Lotfollahi, P. (2022) One new species and two new records for Iran from the mites of family Eriophyidae. *24th Iranian Plant Protection Congress, 3-6 September, IRIPP, Tehran*, pp. 450–451.
- Tarasi, J. & Taghaddosi, M.V. (2005) Record of *Anthocoptes salicis* (Acari: Eriophyidae) from Iran. *Journal of Entomological Society of Iran*, 25(1): 77–78.
- The World Flora Online (2025) World Flora Online. Available from: <http://www.worldfloraonline.org/taxon/wfo-0000125895> (Accessed on 01 August 2025).