



Persian J. Acarol., 2024, Vol. 13, No.3, pp. 361–373.
https://doi.org/10.22073/pja.v13i3.85509
Journal homepage: <http://www.biotaxa.org/pja>



<http://zoobank.org/urn:lsid:zoobank.org:pub:B5F1694D-3F70-481B-AD6C-997D6F6C132B>

Article

Additions to the oribatid mite (Acari, Oribatida) fauna of Guatemala, with description of a new species of *Pergalumna*

Sergey G. Ermilov 

Tyumen State University, Institute of Environmental and Agricultural Biology (X-BIO), Tyumen, Russia; E-mail: ermilovacari@yandex.ru

ABSTRACT

The present study is based on oribatid mite material collected from five locations in Eastern Guatemala. A list of 98 species, belonging to 68 genera and 43 families, is presented; 70 species are recorded for the first time in Guatemalan fauna. A new species of genus *Pergalumna*, *P. paraobsidiana* **sp. nov.** is described from rainforest litter.

KEY WORDS: Central America, fauna, galumnid mites, morphology, new record, new species, taxonomy.

PAPER INFO.: Received: 15 April 2024, Accepted by: *O. Joharchi*, 1 May 2024, Published: 15 July 2024

INTRODUCTION

Oribatid mites (Acari, Oribatida) of Guatemala are insufficiently studied (Stoll 1893; Willmann 1930; Woolley 1966; Hammer 1969; Mahunka 1978, 1983; Behan-Pelletier and Ryabinin 1991; Schatz 2006). The main goals of our paper are as follows: to present a list of all identified taxa (including new records) collected from five locations in Eastern Guatemala, and to describe one new species belonging to the nominate subgenus of *Pergalumna* Grandjean, 1936 (Galumnidae) as *Pergalumna paraobsidiana* **sp. nov.**

The genus *Pergalumna* was proposed by Grandjean (1936), with *Oribata nervosa* Berlese, 1914 as type species. The nominate subgenus comprises more than 180 species having a cosmopolitan distribution collectively (Subías 2022, online version 2024). The generic and subgeneric diagnoses were revised by Ermilov and Klimov (2017). An identification key to the Neotropical species of *Pergalumna* has been presented by Ermilov and Starý (2021).

Prior this study, two species of Galumnidae have been reported in the oribatid fauna of Guatemala, *Allogalumna microporosa* Mahunka, 1978 and *Neopilizetes rugifrons* (Stoll, 1893).

MATERIAL AND METHODS

Material – Samples were collected from five locations in Eastern Guatemala: 1 – Zacapa Department, 6.3 km NE Highway, sifting dry leaf litter, 8.VII.1986 (J.M. Campbell); 2 – Zacapa Department, San Lorenzo 4 mi. N, sifted mushroom piles, 18.VII.1986 (J.M. Campbell); 3 – Zacapa Department, Sierra de Las Minas, San Lorenzo, 1700 m a.s.l., lichens and moss mats on rocks, 7–10.XI.1986 (E.E. Lindquist); 4 – Zacapa Department, Sierra de Las Minas, 9 km N. San Lorenzo,

How to cite: Ermilov, S.G. (2024) Additions to the oribatid mite (Acari, Oribatida) fauna of Guatemala, with description of a new species of *Pergalumna*. *Persian Journal of Acarology*, 13(3): 361–373.

2100 m a.s.l., damp litter in cloud forest, 17.XI.1986 (E.E. Lindquist); 5 – Izabal Department, Las Escobar, 8 km SW Puerto Barrios, rainforest litter, 12–14.XI.1986 (E.E. Lindquist).

Observation and documentation – For measurement and illustration, specimens were mounted in lactic acid on temporary cavity slides. All measurements are in micrometers (μm); body length was measured in lateral view, from the tip of the rostrum to the posterior edge of the notogaster; notogastral width refers to the maximum width in dorsal aspect (behind pteromorphs); setal lengths were measured perpendicular to their long axes, accounting for curvature. Formulas for leg setation are given in parentheses according to the sequence trochanter-femur-genu-tibia-tarsus (famulus included); formulas for leg solenidia are given in square brackets, according to the sequence genu-tibia-tarsus. Drawings were made with a camera lucida using a Leica DM 2500 transmission light microscope.

Terminology – Morphological terminology used in this paper mostly follows that of Grandjean (1956, 1957), Norton and Ermilov (2017) and Ermilov *et al.* (2023); also, Ermilov and Klimov (2017) for overview.

Abbreviations – The following morphological abbreviations are used: *Prodorsum*: *L* = lamellar line; *S* = sublamellar line; *N* = prodorsal leg niche; *E*, *T* = lateral ridges; *ro*, *le*, *in*, *bs* = rostral, lamellar, interlamellar, and bothridial setae, respectively; *Ad* = dorsosejugal porose area; *D* = dorsophragma; *P* = pleurophragma. *Notogaster*: *Aa*, *A1*, *A3* = porose areas; *c*, *la*, *lm*, *lp*, *h*, *p* = setae; *ia*, *im*, *ip*, *ih*, *ips* = lyrifissures; *gla* = opisthonotal gland opening. *Gnathosoma*: *a*, *m*, *h* = subcapitular setae; *or* = adoral seta; *sup*, *inf*, *d*, *l*, *cm*, *acm*, *ul*, *su*, *vt*, *lt* = palp setae; ω = palp solenidium; *as* = axillary saccule; *cha*, *chb* = cheliceral setae; *Tg* = Trägårdh's organ. *Epimeral and lateral podosomal regions*: *1a*, *3b*, *3c*, *4a*, *4b*, *4c* = epimeral setae; *PdI*, *PdII* = pedotecta I, II, respectively; *dis* = discidium; *cir* = circumpedal carina. *Anogenital region*: *g*, *ag*, *an*, *ad* = genital, aggenital, anal, and adanal setae, respectively; *iad* = adanal lyrifissure; *po* = preanal organ. *Legs*: *Tr*, *Fe*, *Ge*, *Ti*, *Ta* = trochanter, femur, genu, tibia, and tarsus, respectively; ω , ϕ , σ = solenidia; ε = famulus; *d*, *l*, *v*, *bv*, *ev*, *ft*, *tc*, *it*, *p*, *u*, *a*, *s*, *pv*, *pl* = setae; *pa* = porose area.

LIST OF TAXA

Distribution mostly from Subías (2022, online version 2024). Ptyctimous mites and some Damaeidae are not included.

Brachychthoniidae

Liochthonius fimbriatissimus Hammer, 1962: location 4 (1 ex.). Distribution: Australasian, Neotropical, Antarctic.

Sphaerochthoniidae

Sphaerochthonius splendidus (Berlese, 1904): location 1 (7 ex.). Distribution: Tropical, Subtropical.

Hypochthoniidae

Eohypochthonius crassisetiger Aoki, 1959: location 5 (2 ex.). Distribution: Southern Palaeartic, Australasian, Neotropical.

Eohypochthonius gracilis (Jacot, 1936): location 5 (1 ex.). Distribution: Tropical, Subtropical.

Epilohmanniidae

Epilohmannia minuta Berlese, 1920: location 5 (2 ex.). Distribution: Tropical, Central and Eastern U.S.A.

Epilohmannoides (Neoepilohmannoides) xena (Mahunka, 1983): location 5 (1 ex.). Distribution: Northern Neotropical, Vietnam, Iran.

Trhypochthoniidae

Allonothrus tuxtlasensis Palacios-Vargas & Iglesias, 1997: location 5 (1 ex.). Distribution: Northern Neotropical.

Archezogetes magnus (Sellnick, 1925): location 5 (47 ex.). Distribution: Tropical, Subtropical.

Trhypochthonius tectorum (Berlese, 1896): location 3 (13 ex.). Distribution: Semicosmopolitan.

Malaconothridae

Malaconothrus aureopunctatus Hammer, 1979: location 4 (2 ex.). Distribution: Oriental, Brazil, Ethiopia.

Nothridae

Nothrus anauniensis Canestrini & Fanzago, 1877: location 3 (1 ex.). Distribution: Cosmopolitan.

Nothrus willmanni Mahunka, 1983: location 4 (7 ex.). Distribution: Neotropical.

Nanhermanniidae

Nanhermannia elegantissima Hammer, 1958: location 4 (4 ex.). Distribution: Neotropical, Subantarctic.

Hermanniellidae

Hermanniella sp.: 4 (7 ex.).

Sacculobates horologiorum Grandjean, 1962: location 5 (4 ex.). Distribution: Neotropical.

Plasmobatidae

Plasmobates cribratus Grandjean, 1929: location 5 (4 ex.). Distribution: Neotropical.

Licnodamaeidae

Licnodamaeus sp.: location 3 (1 ex.).

Pheroliodidae

Pheroliodes wehnckeii (Willmann, 1930): location 3 (3 ex.). Distribution: Neotropical.

Gymnodamaeidae

Jacotella ornata (Balogh & Csiszár, 1963): location 3 (2 ex.). Distribution: Neotropical, India.

Damaeidae

Epidamaeus sp.: location 3 (1 ex.).

Cepheusidae

Oribatodes mirabilis Banks, 1895: location 4 (1 ex.). Distribution: Northern Nearctic.

Polypterozetidae

Polypterozetes cherubin Berlese 1916: location 4 (9 ex.). Distribution: Italy.

Basilobelbidae

Basilobelba wernerii Mahunka, 1982: locations 4 (18 ex.), 5 (4 ex.). Distribution: Northern

Neotropical.

Eremaeidae

Eueremaeus sp.: location 3 (1 ex.).

Astegistidae

Cultroribula bicuspidata Mahunka, 1978: location 5 (5 ex.). Distribution: Tropical, Subtropical.

Furcoppia sp.: location 5 (2 ex.).

Peloppiidae

Ceratorchestes cf. *baloghi* Mahunka, 1983: locations 1 (1 ex.), 5 (1 ex.). Distribution: Northern Neotropical.

Metrioppia serrata (Sellnick, 1931): location 4 (14 ex.). Distribution: Mexico, U.S.A. (Virginia).

Eremulidae

Eremulus brasiliensis Pérez-Íñigo & Baggio, 1985: locations 1 (2 ex.), 3 (5 ex.), 4 (9 ex.), 5 (5 ex.). Distribution: Neotropical.

Damaeolidae

Fosseremus laciniatus (Berlese, 1905): location 4 (1 ex.). Distribution: Cosmopolitan.

Oppiidae

Amboroppia (*Quintanoppia*) *guatemalaensis* Ermilov, 2024: location 4 (7 ex.). Distribution: Guatemala.

Amboroppia (*Quintanoppia*) *zacapaensis* Ermilov, 2024: location 4 (4 ex.). Distribution: Guatemala.

Arcoppia dechambrierorum (Mahunka, 1983): location 5 (9 ex.). Distribution: Northern Neotropical.

Arcoppia tumulosa Ermilov, 2024: locations 1 (1 ex.), 4 (4 ex.). Distribution: Guatemala.

Arcoppia sp.: 5 (12 ex.).

Mimoppia (*Dysarcoppia*) *dendropectinata* (Woas, 1986): location 5 (18 ex.). Distribution: El Salvador.

Neoamerioppia sp.: location 4 (1 ex.).

Oppiella nova (Oudemans, 1902): locations 1 (1 ex.), 4 (2 ex.). Distribution: Cosmopolitan.

Pararamusella cf. *disjuncta* Mahunka & Palacios-Vargas, 1998: location 4 (16 ex.). Distribution: Mexico. Remarks. Our specimens are similar to those from the original description (Mahunka & Palacios-Vargas, 1998), but they have narrowed (nearly pointed) rostrum.

Pseudoamerioppia barrancensis (Hammer, 1961): location 1 (18 ex.). Distribution: Neotropical, Philippines, western Africa, Canary Islands.

Ramusella (*Ramusella*) *chulumaniensis* (Hammer, 1958): locations 4 (1 ex.), 5 (2 ex.). Distribution: Neotropical, Oriental.

Ramusella (*Insculptoppia*) *claudelionsi* (Călugăr & Vasiliu, 1983): location 4 (1 ex.). Distribution: Cuba.

Wallworkoppia cervifer (Mahunka, 1983): locations 4 (1 ex.), 5 (3 ex.). Distribution: Northern Neotropical.

Sternoppiidae

Sternoppia incisa Balogh & Mahunka, 1977: locations 4 (5 ex.), 5 (9 ex.). Distribution: Neotropical.

Quadropiidae

Quadropia (*Coronoquadropia*) *circumita* (Hammer, 1961): location 5 (1 ex.). Distribution: Tropical, Subtropical.

Suctobelbidae

Allosuctobelba sp.: location 4 (3 ex.).

Suctobelbella (*Suctobelbella*) *subcornigera maculata* Shtanchaeva & Subías, 2009: locations 4 (1 ex.), 5 (1 ex.). Distribution: Holarctic, Nearctic, Argentina.

Suctobelbella (*Suctobelbella*) *perdentata* (Hammer, 1961): location 5 (1 ex.). Distribution: Neotropical.

Suctobelbella (*Flagrosuctobelba*) *crassisetosa* Ermilov, 2024: location 5 (1 ex.). Distribution: Mexico.

Suctobelbella (*Flagrosuctobelba*) *forsslundi* (Strenzke, 1950): location 4 (3 ex.). Distribution: Palaearctic, Cuba.

Suctobelbilla *tetrasetosa* Ermilov, 2024: location 5 (1 ex.). Distribution: Mexico.

Dampfiellidae

Beckiella sp.: location 4 (1 ex.).

Otocephidae

Cavernocephus longicristatus Ermilov, 2024: location 4 (9 ex.). Distribution: Guatemala.

Cavernocephus undulatus P. Balogh, 2002: location 5 (1 ex.). Distribution: Central America.

Carabodidae

Carabodes (*Carabodes*) *guatemalaensis* Ermilov, 2023: locations 1 (3 ex.), 4 (3 ex.). Distribution: Guatemala.

Carabodes (*Klapperiches*) *lindquisti* Ermilov, 2023: location 3 (9 ex.). Distribution: Guatemala.

Carabodes (*Klapperiches*) *venezolanus* Subías & Arillo, 2004: locations 3 (17 ex.), 5 (1 ex.). Distribution: Neotropical.

Costacarabodes turrialbai Fernández, Theron, Leiva & Jordaan, 2018: location 5 (4 ex.). Distribution: Neotropical.

Gymnobodes minimus Ermilov, 2016: location 5 (3 ex.). Distribution: Cuba.

Tectocephidae

Tectocephus sarekensis Trägårdh, 1910: locations 1 (2 ex.), 4 (4 ex.). Distribution: Cosmopolitan.

Tectocephus velatus (Michael, 1880): locations 2 (1 ex.), 4 (1 ex.). Distribution: Cosmopolitan.

Microtegeidae

Microtegeus borhidii Balogh & Mahunka, 1974: locations 1 (1 ex.), 4 (2 ex.), 5 (6 ex.). Distribution: Neotropical, Oriental, Japan.

Phenopelopidae

Eupelops acromios (Hermann, 1804): location 5 (1 ex.). Distribution: Semicosmopolitan.

Microzetidae

Protozetes janosbaloghi Ermilov, 2024: locations 4 (3 ex.), 5 (4 ex.). Distribution: Guatemala.

Schalleria mexicana Mahunka & Palacios-Vargas, 1996: location 5 (1 ex.). Distribution: Mexico.

Epactozetidae

Truncozetes sp.: location 5 (5 ex.).

Oribatellidae

Oribatella dechambrieri Mahunka, 1983: location 4 (6 ex.). Distribution: Northern Neotropical.

Oribatella sp. A: location 4 (1 ex.).

Oribatella sp. B: location 5 (15 ex.).

Oribatella sp. C: location 5 (3 ex.).

Ceratozetidae

Ceratozetes mediocris Berlese, 1908: location 4 (4 ex.). Distribution: Semicosmopolitan.

Fuscozetes fuscipes (Koch, 1844): location 1 (6 ex.). Distribution: Holarctic, Costa Rica, Vietnam.

Punctoribatidae

Lamellobates molecula (Berlese, 1916): locations 1 (44 ex.), 2 (1 ex.). Distribution: Tropical, Subtropical.

Mochlozetidae

Dynatozetes magnus (Banks, 1895): locations 4 (3 ex.), 5 (3 ex.). Distribution: Nearctic, Ecuador.

Scheloribatidae

Scheloribates cf. *elegans* Hammer, 1958: locations 1 (4 ex.), 3 (5 ex.), 4 (19 ex.). Distribution: Neotropical, Oriental, Hawaii.

Scheloribates laevigatus (Koch, 1835): location 1 (1 ex.). Distribution: Semicosmopolitan.

Scheloribates pallidulus (Koch, 1841): locations 1 (1 ex.), 5 (7 ex.). Distribution: Semicosmopolitan.

Scheloribates praeincisus (Berlese, 1910): locations 1 (6 ex.), 2 (1 ex.), 3 (1 ex.). Distribution: Oriental, Australasian, Afrotropical, Neotropical, Southern Holarctic.

Scheloribates (*Bischeloribates*) sp.: location 3 (1 ex.).

Scheloribates (*Perscheloribates*) *mexicoensis* Ermilov & Kontschán, 2023: location 5 (1 ex.). Distribution: Mexico.

Scheloribates (*Perscheloribates*) *minusculus* Hammer, 1961: locations 3 (1 ex.), 5 (1 ex.). Distribution: Neotropical.

Scheloribates (*Topobates*) sp.: location 5 (2 ex.).

Oripodidae

Oripoda elongata Banks & Pergande, 1904: location 3 (1 ex.). Distribution: U.S.A., Mexico.

Haplozetidae

Peloribates longisetosus (Willmann, 1930): locations 1 (2 ex.), 3 (10 ex.). Distribution: Neotropical, eastern Palaearctic, India.

Protoribates capucinus Berlese, 1908: location 5 (3 ex.). Distribution: Cosmopolitan.

Protoribates paracapucinus (Mahunka, 1988): location 1 (1 ex.). Distribution: Tropical and Subtropical.

Protoribates sp.: location 4 (5 ex.).

Setoxylobates palaciosvargasi Ermilov & Kontschán, 2023: location 5 (21 ex.). Distribution: Mexico.

Rostrozetes ovulum (Berlese, 1908): locations 1 (6 ex.), 4 (1 ex.), 5 (9 ex.). Distribution: Tropical and Subtropical.

Galumnidae

Galumna flabellifera Hammer, 1958: location 1 (6 ex.). Distribution: Tropical and Subtropical.

Galumna laselvae P. Balogh, 1997: location 1 (1 ex.). Distribution: Neotropical.

Galumna sp.: location 1 (1 ex.).

Pergalumna horvathorum P. Balogh, 1997: location 5 (12 ex.). Distribution: Northern Neotropical.

Pergalumna paraboliviana Ermilov & Gwiazdowicz, 2015: location 4 (39 ex.). Distribution: Peru.

Pergalumna paraobsidiana **sp. nov.**: location 5 (6 ex.).

Pergalumna sp.: locations 1 (1 ex.), 4 (3 ex.).

Trichogalumna nipponica (Aoki, 1966): location 4 (3 ex.). Distribution: Semicosmopolitan.

Galumnellidae

Galumnopsis holoscripta Grandjean, 1931: location 5 (2 ex.). Distribution: Neotropical.

The list includes 98 species belonging to 68 genera and 43 families. Of the 79 species (except one new species and 18 unidentified species), seven species are known only from Guatemala, one is Nearctic, one is European, 31 are Neotropical, and 27 (including 12 Cosmopolitan/Semicosmopolitan species) have broad distributions (more than one geographical region). Except *Nothrus willmanni*, *Sacculobates horologiorum*, *Peloribates longisetosus*, 18 unidentified species and the seven Guatemalan species, the other species (70) are recorded for the first time in Guatemala.

TAXONOMY

Family Galumnidae

Pergalumna Grandjean, 1936

Subgenus *Pergalumna* (*Pergalumna*) Grandjean, 1936

Type species: *Oribata nervosa* Berlese, 1914

Pergalumna paraobsidiana sp. nov. (Figs. 1–12)

<http://zoobank.org/urn:lsid:zoobank.org:act:2128D02D-CAA8-4DEF-8CF8-7AECB8779CF7>

Type material

Holotype (male) and five paratypes (two males and three females): Eastern Guatemala, Izabal Department, Las Escobar, 8 km SW Puerto Barrios, rainforest litter, 12–14.XI.1986 (E.E. Lindquist).

Type deposition

The holotype is deposited in the collection of the Canadian National Collection, Ottawa, Canada; five paratypes are in the collection of the Tyumen State University Museum of Zoology, Tyumen, Russia. All specimens are preserved in 70% solution of ethanol with a drop of glycerol.

Diagnosis

Body length: 525–570. Prodorsum, pteromorph, ventral plate, and posterior part of notogaster densely tuberculate; anterior and central parts of notogaster with dense short longitudinal ridges. Rostrum pointed. Rostral, lamellar and interlamellar setae long, setiform, barbed; relative length: $in > le > ro$; bothridial seta long, setiform, unilaterally shortly ciliate. Dorsosejugal porose area present.

Dorsosejugal suture present but indistinct. Three pairs of porose areas present: *Aa* and *A3* oval; *A1* elongate oval; *Aa* located between *la* and *lm*, closer and anteromedially to *la*. Median pore absent. Circumpedal carina directed to insertion of seta *3b*. Epimeral and anogenital setae comparatively short. Postanal porose area absent. Leg famulus curved mediolaterally, directed forwards, inserted between solenidia ω_1 and ω_2 ; solenidion on tibia IV with well-developed swollen mediobasal part, inserted in posterior part of the segment.

Description

Measurements – Body length: 540 (holotype), 525–570 (paratypes); notogaster width: 405 (holotype), 375–420 (paratypes). No differences between males and females in body size.

Integument – Body color brown. Body surface densely microgranulate sculpturing; additionally, prodorsum, pteromorph, ventral plate (including subcapitular mentum and genital and anal plates), and posterior part of notogaster densely tuberculate; anterior and central parts of notogaster with dense short longitudinal ridges.

Prodorsum – Rostrum pointed. Lamellar and sublamellar lines thin, parallel, curving backwards mediolaterally. Rostral (67–75), lamellar (97–105) and interlamellar (124–127) setae setiform, barbed; *le* removed from *L*; bothridial seta (153–161) setiform, unilaterally shortly ciliate. Dorsosejugal porose area elongate oval (15–19 × 5), located posterolaterally to insertion of *in*. Dorsophragma comparatively long.


Notogaster – Dorsosejugal suture present but indistinct. All notogastral setae represented by setal alveoli. Three pairs of porose areas developed; *Aa* (30–37 × 15–19) and *A3* (22–30 × 15–19) oval; *A1* (45–60 × 15–19) elongate oval; *Aa* located between *la* and *lm*, closer and anteromedially to *la*. Median pore absent. Opisthotal gland opening and all lyrifissures distinct: *gla* located laterally to *A1*; *im* anterolaterally to *A1*; *ip* between *p*₁ and *p*₂, closer to *p*₂; *ih* and *ips* close to each other, anteriorly to *p*₃.

Gnathosoma – Size of subcapitulum: 139–142 × 120–124; subcapitular (*a*, *h*: 28–30; *m*: 30–34) and adoral (15–17) setae setiform, barbed. Length of palp: 105–112; setation: 0–2–1–3–9(+ ω); postpalpal seta (5) spiniform, smooth. Length of chelicera: 187–191; setae (*cha*: 52–56; *chb*: 37–39) setiform, barbed.

Epimeral and lateral podosomal regions – Epimeral setal formula: 1–0–2–3; setae (*4a*, *4b*: 11; *1b*: 17–19; *3b*: 22–26; *3c*, *4c*: 30–34) setiform, smooth. Circumpedal carina medium-sized, directed to insertion of seta *3b*.

Anogenital region – Anogenital formula: 6–1–2–3; genital (*g*₁: 17–19; others: 11), aggenital (11), anal (11), and adanal (11) setae setiform, roughened; anterior edge of genital plate with two setae; aggenital seta located between genital and anal apertures, slightly closer to the former; adanal setae *ad*₁ and *ad*₂ posteriorly, *ad*₃ laterally to anal plate; distance *ad*₁–*ad*₂ shorter than *ad*₂–*ad*₃. Adanal lyrifissure located close and parallel to anal plate. Postanal porose area absent.

Legs – Median claw distinctly thicker than lateral claws; all claws slightly barbed on dorsal side. Porose area on femora I–IV and on trochanters III, IV poorly visible; proximoventral porose area on tarsi and distoventral porose area on tibiae not observable. Formulas of leg setation and solenidia: I (1–4–3–4–20) [1–2–2], II (1–4–3–4–15) [1–1–2], III (1–2–1–3–15) [1–1–0], IV (1–2–2–3–12) [0–1–0]; homology of setae and solenidia indicated in Table 1; famulus curved mediolaterally, directed forwards, inserted between solenidia ω_1 and ω_2 ; solenidion on tibia IV with well-developed swollen mediobasal part, inserted in posterior part of the segment.




Figures 1–4. *Pergalumna paraobsidiana* sp. nov. (adult) – 1. Dorsal view of body; 2. Ventral view of body (legs and right pteromorph not shown); 3. Right lateral view of body (gnathosoma, legs and right pteromorph not shown); 4. Posterior view of body (part of left half not shown).

Table 1. Leg setation and solenidia of adult *Pergalumna paraobsidiana* sp. nov.

Leg	Tr	Fe	Ge	Ti	Ta
I	v'	d, (l), bv''	(l), v', σ	(l), (v), φ ₁ , φ ₂	(ft), (tc), (it), (p), (u), (a), s, (pv), v', (pl), l'', ε, ω ₁ , ω ₂
II	v'	d, (l), bv''	(l), v', σ	(l), (v), φ	(ft), (tc), (it), (p), (u), (a), s, (pv), ω ₁ , ω ₂
III	v'	d, ev'	l', σ	l', (v), φ	(ft), (tc), (it), (p), (u), (a), s, (pv)
IV	v'	d, ev'	d, l'	l', (v), φ	ft'', (tc), (p), (u), (a), s, (pv)

Note: Roman letters refer to normal setae; Greek letters to solenidia (except ε = famulus); single quotation mark (') designates setae on the anterior and double quotation mark (') setae on the posterior side of a given leg segment; parentheses refer to a pair of setae.



Figures 5–12. *Pergalumna paraobsidiana* sp. nov. (adult) – 5. Anterior view of prodorsum, dorsoanterior view; 6. Subcapitulum, ventral view; 7. Palp, left, paraxial view; 8. Chelicera, left, paraxial view; 9. Leg I, right, antiaxial view; 10. Leg II (trochanter not shown), right, antiaxial view; 11. Leg III, left, antiaxial view; 12. Leg IV, left, antiaxial view.

Comparison

Pergalumna obsidiana sp. nov. is most similar to *P. obsidiana* Palacios-Vargas & Villagómez, 2017 and *P. ekaterinae* Páez, Villagómez & Palacios-Vargas, 2019 from Mexico in the following main traits: rostrum pointed; bothridial seta long, setiform, shortly ciliate; rostral, lamellar and

interlamellar setae long; solenidion of the leg tibia IV with mediobasal swollen part.

The new species differs from *P. obsidiana* by the following traits: body size larger (length: 525–570 versus 443–513); anterior and central parts of notogaster with dense short longitudinal ridges (versus tuberculate); genital plate tuberculate (versus with strong transverse ridges); and rostral region without longitudinal ridge (versus with ridge).

The new species differs from *P. ekaterinae* by the following traits: body size larger (length: 525–570 versus 434–503); prodorsum, pteromorph, ventral plate, and posterior part of notogaster densely tuberculate, and anterior and central parts of notogaster with dense short longitudinal ridges (versus surface without heavy sculpturing); notogaster with three pairs (versus four pairs) of notogastral porose areas; posterior part of notogaster without striate bands (versus with bands).

Etymology

The species name *paraobsidiana* refers to the similarity between the new species and *P. obsidiana*.

ACKNOWLEDGEMENTS

I cordially thank J.M. Campbell and E.E. Lindquist for collecting the oribatid materials, W. Knee for facilitating the loan of Guatemalan Acari material from the Canadian National Collection of Insects, Arachnids and Nematodes (Ottawa, Canada) and two anonymous reviewers for valuable comments.

REFERENCES

- Behan-Pelletier, V.M. & Ryabinin, N.A. (1991) Description of *Sacculozetes filosus* gen. nov., sp. nov. and *Guatemalozetes danos* sp. nov. (Acari: Oribatida) from grassland habitats. *The Canadian Entomologist*, 123: 1135–1147. DOI: [10.4039/Ent1231135-5](https://doi.org/10.4039/Ent1231135-5)
- Berlese, A. (1914) Acari nuovi. Manipulus IX. *Redia*, 10: 113–150.
- Ermilov, S.G. & Klimov, P.B. (2017) Generic revision of the large-winged mite superfamily Galumnoidea (Acari, Oribatida) of the world. *Zootaxa*, 4357: 1–72. DOI: [10.11646/zootaxa.4357.1.1](https://doi.org/10.11646/zootaxa.4357.1.1)
- Ermilov, S.G., Kolesnikov, V.B., Kontschán, J. & Klimov, P.B. (2023) Taxonomic contribution to the knowledge of Galumnidae (Acari, Oribatida, Galumnidae) from Cuba. *Zootaxa*, 5258: 465–474. DOI: [10.11646/zootaxa.5258.4.7](https://doi.org/10.11646/zootaxa.5258.4.7)
- Ermilov, S.G. & Starý, J. (2021) A new species of *Pergalumna* (Acari, Oribatida, Galumnidae) from Bolivia, with updated identification key to known species of the genus from the Neotropical region. *Systematic and Applied Acarology*, 26: 1097–1108. DOI: [10.11158/saa.26.6.7](https://doi.org/10.11158/saa.26.6.7)
- Grandjean, F. (1936) Les Oribates de Jean Frédéric Hermann et de son père. *Annales de la Société Entomologique de France*, 105: 27–110.
- Grandjean, F. (1956) Galumnidae sans carènes lamellaires (Acariens, Oribates). 1^{re} série. *Bulletin de la Société Zoologique de France*, 81: 134–150.
- Grandjean, F. (1957) Galumnidae sans carènes lamellaires (Acariens, Oribates). 2^e série. *Bulletin de la Société Zoologique de France*, 82: 57–71.
- Hammer, M. (1969) Oribatids found at plant quarantine stations in U.S.A. *Videnskabelige Meddelelser fra Dansk naturhistorisk Forening i Kjøbenhavn*, 132: 63–78.
- Mahunka, S. (1978) Neue und interessante Milben aus dem Genfer Museum XXV. On some oribatids collected by Dr. P. Strinati in Guatemala (Acari: Oribatida). *Acarologia*, 20: 133–142.
- Mahunka, S. (1983) Neue und interessante Milben aus dem Genfer Museum XLVII. Oribatida

- Americana 7: Guatemala II. *Revue suisse de Zoologie*, 90: 709–724.
- Mahunka, S. & Palacios-Vargas, J.G. (1998) New oppiid oribatid mites from Mexico (Acari: Oribatida) I. *Acta Zoologica Academiae Scientiarum Hungaricae*, 44: 283–295.
- Norton, R.A. & Ermilov, S.G. (2017) Identity of the oribatid mite *Oribata curva* and transfer to *Trichogalumna* (Acari, Oribatida, Galumnidae), with discussion of nomenclatural and biogeographical issues in the ‘*curva*’ species-group. *Zootaxa*, 4272: 551–564. DOI: [10.11646/zootaxa.4272.4.4](https://doi.org/10.11646/zootaxa.4272.4.4)
- Páez, J., Villagomez, F. & Palacios-Vargas, J.G. (2019) Description of a new *Pergalumna* (Acari: Oribatida: Galumnidae) species from Mexico and its postembryonic development. *Zootaxa*, 4647: 385–406. DOI: [10.11646/zootaxa.4647.1.25](https://doi.org/10.11646/zootaxa.4647.1.25)
- Palacios-Vargas, J.G. & Villagomez, F. (2017) Three new species of *Pergalumna* (Acari: Oribatida: Galumnidae) from the tropical rainforest of Veracruz, Mexico. *Zootaxa*, 4242: 77–94. DOI: [10.11646/zootaxa.4242.1.4](https://doi.org/10.11646/zootaxa.4242.1.4)
- Schatz, H. (2006) Catalogue of known oribatid mite species (Acari Oribatida) from the Central American landbridge (First part). *Tropical Zoology*, 19: 209–288.
- Stoll, O. (1893) *Arachnida Acaridea. Biologia Centrali-Americana*. In: Godman, F.D. & Salvin, O. (Eds.), *Biologia Centrali-Americana*. Privately printed (1886–1893) by R. H. Porter, London. pp. i–xxi, 1–55.
- Subías, L.S. (2022) Listado sistemático, sinonímico y biogeográfico de los ácaros oribátidos (Acariformes, Oribatida) del mundo (excepto fósiles). *Monografías Electrónicas Sociedad Entomológica Aragonesa*, 12: 1–539.
- Subías, L.S. (2024) Listado sistemático, sinonímico y biogeográfico de los ácaros oribátidos (Acariformes: Oribatida) del mundo (excepto fósiles), 19ª actualización, pp. 1–545. Available from: http://bba.bioucm.es/cont/docs/RO_1.pdf (Accessed in January 2024).
- Willmann, C. (1930) Neue Oribatiden aus Guatemala. *Zoologischer Anzeiger*, 88: 239–246.
- Woolley, T.A. (1966) New oribatid mites from Central America (Acari: Cryptostigmata). *Proceedings of the Entomological Society of Washington*, 68: 334–339.

COPYRIGHT

Ermilov. Persian Journal of Acarology is under a free license. This open-access article is distributed under the terms of the Creative Commons-BY-NC-ND which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original author and source are credited.

موارد اضافه شده به فون هرناهای اورibatید (Acari, Oribatida) گواتمالا، همراه با توصیف گونه جدیدی از *Pergalumna*

سرگی جی. ارمیلوف

دانشگاه ایالتی تیومن، مؤسسه زیست‌شناسی زیست‌محیطی و کشاورزی (X-BIO)، تیومن، روسیه؛ رایانامه: ermilovacari@yandex.ru

چکیده

مطالعه حاضر بر اساس نمونه‌های هرناهای اورibatید جمع‌آوری شده از پنج مکان در شرق گواتمالا است. فهرستی از ۹۸ گونه، متعلق به ۶۸ جنس و ۴۳ خانواده، ارائه شده است؛ ۷۰ گونه برای نخستین بار برای فون گواتمالا ثبت شده است. گونه جدیدی از جنس *P. Pergalumna*، *paraobsidiana* sp. nov. از بستر جنگل‌های بارانی توصیف می‌شود.

واژگان کلیدی: آمریکای مرکزی، فون، کنه‌های گالومنید، ریخت‌شناسی، گزارش جدید، گونه جدید، آرایه‌شناسی.

اطلاعات مقاله: تاریخ دریافت: ۱۴۰۳/۱/۲۷، تاریخ پذیرش توسط / جوهرچی: ۱۴۰۳/۲/۲۲، تاریخ چاپ: ۱۴۰۳/۴/۲۵