



*Persian J. Acarol.*, 2020, Vol. 9, No. 3, pp. 243–253.  
<http://dx.doi.org/10.22073/pja.v9i3.60194>  
Journal homepage: <http://www.biotaxa.org/pja>



## Article

### New records and remarks on Tydeoidea (Acari: Trombidiformes) from Mazandaran province of Iran

Maryam Darbemamieh<sup>1\*</sup>, Andrzej Kaźmierski<sup>2</sup> and Saeid Paktinat-Saeij<sup>3</sup>

1. Department of Plant Protection, Faculty of Agriculture, Campus of Agriculture and Natural Resources, Razi University, Kermanshah, Iran; E-mail: [m.darbemamieh@razi.ac.ir](mailto:m.darbemamieh@razi.ac.ir)
2. Department of Animal Morphology, Faculty of Biology, Adam Mickiewicz University, Umultowska 89, 61-614 Poznan, Poland; E-mail: [aKaźmierski@tlen.pl](mailto:aKaźmierski@tlen.pl)
3. Department of Plant Protection, Faculty of Crop Sciences, Sari Agricultural Sciences and Natural Resources University, Sari, Iran; E-mail: [saeedpaktinat@yahoo.com](mailto:saeedpaktinat@yahoo.com)

\* Corresponding author

#### ABSTRACT

Two collections of Tydeoidea from Mazandaran province revealed 16 species from six genera in two families. Of these, *Tydeus lindquisti* (Marshall, 1970) is a new record for Iran and 12 species are new records for the province. Photographs and supplementary data are provided for some species.

**KEY WORDS:** *Homeopronematus*; Iolinidae; *Lorryia*; *Neopronematus*; *Proctotydaeus*; *Pronematus*; *Tydeus lindquisti*.

**PAPER INFO.:** Received: 9 March 2020, Accepted: 23 April 2020, Published: 15 July 2020

#### INTRODUCTION

The superfamily Tydeoidea Kramer, 1877 is found worldwide and contains over 700 species and four families: Triophtydeidae André, 1980, Ereyneidae Oudemans, 1931, Tydeidae Kramer, 1877 and Iolinidae Pritchard, 1956 (André and Fain 2000; Zhang *et al.* 2011; Darbemamieh *et al.* 2015; Silva *et al.* 2016). Although these mites are numerous and common in soil, litter or moss, and on plants, information about their distribution, biology, feeding specialization and zoogeography is rudimentary (Kaźmierski 2008). On the other hand, some arboreal tydeoids have an important role in providing alternative food for phytoseiid mites in absence of other prey (Flaherty and Hoy 1971; Calvert and Huffaker 1974; Niemczyk and Kaźmierski, 2002). Some others are scavengers on the leaf surface and may have an important role in health of ecosystems and decreasing the effects of other pests or in the biological control of plant pathogens (English-Loeb *et al.*, 1999). Herbivory is also known for *Tydeus californicus* (Fleschner and Arakawa, 1953). However, some species are also associated with insects (Treat 1970), are predators of eriophyids (Ahmad-Hosseini *et al.* 2019) or are harmful to human and livestock (Kaźmierski 1998).

According to checklist of Kamali *et al.* (2001) and Sadeghi *et al.* (2012), 13 tydeoid species were previously recorded from Mazandaran province including *Pausia magdalenae* (Baker & Delfinado,

**How to cite:** Darbemamieh, M., Kaźmierski, A. & Paktinat-Saeij, S. (2020) New records and remarks on Tydeoidea (Acari: Trombidiformes) from Mazandaran province of Iran. *Persian Journal of Acarology*, 9(3): 243–253.

1976) from citrus, *Triophtydeus immanis* Kuznetzov, 1972 from citrus, tea and pine, *Pronematus sextoni* Baker, 1968, *Lorryia formosa* Cooreman, 1958 from citrus, *L. longina* (Kaźmierski, 1981) from *Cedrus libani* Barrel, *Lorryia obstinata* (Livshitz, 1973) from citrus, *L. oregonensis* (Baker, 1970) from tea and pine, *L. reticulata* (Oudemans, 1928), *Neoapolorryia hippocastani* Kaźmierski, 1998, *Tydeus calabrus* (Castagnoli, 1984) and *Tydeus californicus* (Banks, 1904), all from citrus, *T. gloveri* (Ashmead, 1879) from unknown plant, and *T. inclutus* Livshitz, 1973 from tea, Austrian pine and oriental arborvitae.

The climate of this province is humid, with rich flora, and would potentially harbor a great diversity of tydeoid mites. Our survey presents the results of an investigation in two previous collections made by first and third authors and adds some new records to the Tydeoidea of Mazandaran and Iran. Additional and specific studies will be helpful to illuminate variations and distribution of these mites in northern Iran as well as their importance in agricultural and forest ecosystems.

## MATERIAL AND METHODS

Foliage mites were collected by beating branches over a white plastic board (with a screen above to prevent unwanted material and insects), then transferred into 75% ethanol by means of a fine brush. Also soil mites were extracted by Tullgren funnel, removed with the aid of a stereomicroscope and transferred into 75% ethanol. Some bigger specimens were macerated in lactic acid for a few days prior to being permanently mounted in Hoyer's medium (Walter and Krantz 2009). The preparations were dried in an oven at 50 °C for two weeks. Specimens were examined by means of a phase-contrast microscope (Olympus® BX 51). The nomenclatural terms and setal notations of the idiosoma follows Kaźmierski (1998); those for appendages follows André (1981). We continue to regard *Lorryia* as the senior synonym of *Brachytydeus* as explained by Mondin *et al.* (2016) based on Kaźmierski (1989) and Kaźmierski and Sikora (2008). The measurements are in micrometers (µm). All specimens were deposited in the Acarological collection at Department of Plant Protection, Razi University, Kermanshah, Iran.

## RESULTS

### Family Tydeidae Kramer, 1877

#### Subfamily Tydeinae André, 1979 sensu Kaźmierski, 1996

#### Genus *Tydeus* Koch, 1835 sensu Kaźmierski, 1989

#### *Tydeus caudatus* (Duges, 1834) sensu Baker, 1970

**Remarks** – Some females had several eggs (up to 10 eggs) inside their body. The taxonomic status and some challenges about this species were explained in Darbemamieh *et al.* (2016b). This species has been collected from some other parts of the country (Darbemamieh *et al.* 2010; Akbari *et al.* 2015) but this is the first record from Mazandaran province.

**Examined material** – Angaetaroud village, Noor, 36° 25' 24" N, 52° 13' 14" E, H: 191 m a.s.l., 20 May 2013, soil, 3♀♀.

#### *Tydeus electus* Kuznetzov, 1973

**Remarks** – In this specimen, setae *h2* and *ps1* were clavate to lanceolate, differing from the original description of this species. Other characters match the specific description except the size of the caudal setae that measured about 40 µm. Only one egg was observed inside the female's body. This is the first record of this species from Mazandaran province.

**Examined material** – Chaloos, 36° 39' 41" N, 51° 25' 31" E, H: 10 m a.s.l., orange garden, soil, 4 August 2018, 1♀.

***Tydeus kochi* Oudemans, 1928**

This is the first record of this species from Mazandaran province.

**Examined material** – Izadshahr, 36° 36' 38" N, 52° 10' 26" E, H: -17 m b.s.l., orchard soil, 20 August 2019, 2♀♀.

***Tydeus calabrus* (Castagnoli, 1984)**

**Examined material** – Rostamrood, 36° 35' 21" N, 52° 05' 45" E, H: -21 m b.s.l., apple bark, 20 August 2019, 1♀.

***Tydeus californicus* (Bank, 1904) sensu Baker, 1970**

**Examined material** – Izadshahr, 36° 36' 38" N, 52° 10' 26" E, H: -17 m, orchard soil, 20 August 2019, 1♀.


***Tydeus lindquisti* (Marshall, 1970)**

**Diagnosis** – Body covered with striation and lacking basket weave ornamentation. Dorsal idiosomal setae similar in shape, finely roughened. Cheliceral stylet shorter than palp tarsus. Dorsal striae broad and widely separated and connected in some places. Less than 20 striae between bothridial setae (bo).

**Remarks** – The examined specimens had 17 or 18 striae between the bothridials.

This is the first record of this species from Iran (Fig. 1). This species was categorized in genus *Lorryia* in its first description (Marshall, 1970) and then Kaźmierski (1998) moved it to genus *Tydeus*.

**Examined material** – Alikia-Soltan region, Amol city, 33° 22' 58.26" N, 52° 18' 17.34" E, H: 252 m a.s.l., 5 May 2013, soil, 1♂.



**Figure 1.** Habitus of *Tydeus lindquisti* (left) and its dorsal idiosomal striae (right).

### Genus *Lorryia* Oudemans, 1925

Synonyms: listed in Kaźmierski (1989), and *Quadrotydeus* Momen & Lundqvist 1996 (synonymy by Kaźmierski 1998), *Brachytydeus* Thor, 1931.

#### *Lorryia maga* (Kuznetsov, 1973)

**Diagnosis** – Venilia-type striation (Kaźmierski 1998). Eupathidia of palps cleft distally; gnathosoma protrudes from anterior part of aspidosoma, with only its base concealed (Fig. 2).

This species was first described as *Venilia maga* (Kuznetsov, 1973). This is the first record of this species from Mazandaran province and the first photograph of its dorsum.

**Examined material** – Babol, 36° 32' 58" N, 52° 39' 30" E, H: –6 m b.s.l., orchard soil, 20 August 2019, 1♀.

#### *Lorryia reticulata* (Oudemans, 1928)

**Remarks** – This species was previously classified and described in genus *Tydeus* by Oudemans (Oudemans, 1928). He transferred it to *Lorryia* himself, one year later (Oudemans, 1929). This species is known from several combinations and synonyms summarized in Kaźmierski (1998) and Silva *et al.* (2016) (Fig. 3).

**Examined material** – Chamestan, Angaetaroud village, Noor city, 36° 25' 24" N, 52° 13' 14" E, H: 191 m a.s.l., moss, 1 August 2013, 2♀.



**Figure 2.** Habitus of *Lorryia maga* female, dorsal view (left) and dorsal idiosomal setae and striations (right).



**Figure 3.** Habitus of *Lorryia reticulata*, dorsal view (left), and dorsal idiosomal reticulations (right).

***Lorryia woolleyi* (Baker, 1968)**

**Remarks** – This species was first described as a *Paralorryia* by Baker (1968) but was later moved to *Lorryia* by Sadeghi Namaghi (1995) and Kaźmierski (1998). All dorsal body setae are serrate and sharp distally (Fig. 4) and the movable chelae are not longer than the distal segment of the palps. Round, tear-shaped and finely striated areas around setae, type of body striae (*Paralorryia* s. str.) and longitudinal striae between *c1* and *d1* as shown in figure 4 (right). This is the first record of this species from Mazandaran province.

**Examined material** – Sari, Paen Hoular, 36° 25' 41" N, 53° 08' 01" E, H: 185m, soil, 21 August 2019, 1♀.

**Family Iolinidae Pritchard, 1956**  
**Subfamily Pronematinae André, 1979 sensu André and Fain, 2000**  
**Genus *Proctotydaeus* Berlese, 1911**

**Type species:** *Proctotydaeus viator* Berlese, 1911

This is the first record of this genus from Mazandaran province.


**Subgenus *Proctotydaeus* (*Oriolella*)**

***Proctotydaeus* (*Oriolella*) sp.**

**Examined material** – Soleiman-Abad, Tonekabon, 36° 48' 13" N, 36° 25' 24" E, H: 30 m a.s.l.

walnut, 3 July 2013, 3♀♀.

**Remarks** – femur IV not divided, but tarsus I setae was broken and thus species identification was impossible.



**Figure 4.** *Lorryia woolleyi*, dorsal idiosoma (left), and striae around setae and between *c1* and *d1* (right).

### *Proctotydaeus (Proctotydulus) oblongus* (Kuznetsov, 1973)

This is the first record of this species from Mazandaran province.

**Examined material** – Soleiman-Abad, Tonekabon, 36° 48' 13" N, 36° 25' 24" E, H: 30 m a.s.l., walnut, 3 July 2013, 1♀.

### **Genus *Neopronematus* Panou, Emmanouel and Kaźmierski, 2000**

This is the first record of this genus from Mazandaran province.

**Type species:** *Neopronematus aegeae* Panou, Emmanouel & Kaźmierski, 2000

### *Neopronematus neglectus* (Kuznetsov, 1972)

This is the first record of this species from Mazandaran province.

**Examined material** – Mahmood-Abad, 36° 36' 58" N, 52° 11' 24" E, H: –24 m b.s.l., apple leaf, 20 August 2019, 3♀♀, Babol, 36° 32' 58" N, 52° 39' 30" E, H: –6 m b.s.l., pear leaf, 20 August 2019, 2♀♀.

### *Neopronematus sepasgariani* Sadeghi, Łaniecka & Kaźmierski, 2012

This is the first record of this species from Mazandaran province.

**Examined material** – Mahmood-Abad, apple leaf, 36° 37' 36" N, 52° 13' 48" E, H: –21 m b.s.l., 20 August 2019, 1♀.

### **Genus *Pronematus* Canestrini, 1886**

**Type species:** *Pronematus bonatii* Canestrini, 1886

#### ***Pronematus ubiquitous* (McGregor, 1932)**

This is the first record of this species from Mazandaran province.

**Examined material** – Soleyman-Abad village, Tonekabon, apple leaf, 31 October 2013, 36° 38' 13" N, 52° 50' 10.06" E, H: 30 m a.s.l., 6♀♀.

#### ***Pronematus rykei* Meyer & Rodrigues, 1966**

This is the first record of this species from Mazandaran province.

**Examined material** – Sari, Paein Hoular, 36° 25' 41" N, 53° 08' 01" E, H: 185 m a.s.l., orchard leaf, 4♀♀.

### **Genus *Homeopronematus* André, 1980**

This is the first record of this genus from Mazandaran province.

#### ***Homeopronematus anconai* (Baker, 1943)**

**Remarks** – Dorsal body are approximately long. For example, *e1* (20 µm) longer than *e1-fl* distance. Tectal setae are longer than plurals on tarsus I. This species was first described as a species of *Pronematus* (Baker, 1943) and then moved to *Homeopronematus* by André (1980). This is the first record of this species from Mazandaran province.

**Examined material** – Sari, Paein Hoular, 36° 25' 41" N, 53° 08' 01" E, H: 185m, soil, 21 August 2019, 1♀.

## **DISCUSSION**

Knowledge of Tydeoidea in Iran is limited to a few provinces and studies (Khanjani and Ueckermann 2003; Darbemamieh *et al.* 2010; Sadeghi *et al.* 2012; Akbari *et al.* 2015; Darbemamieh *et al.* 2015, 2016a, b). However, the global knowledge of this group is also in an elementary status. Several factors make them unappealing to study: their delicate bodies and small sizes make them difficult to clarify and mount; many old papers are difficult to find and have inadequate illustrations and descriptions for identification; some descriptions have small figures without enough details or photography; there is no complete key for generic and specific identification in some cases. Collection of these mites is also time-consuming and rarely successful if particular species are being targeted. Most specimens are collected occasionally and by specialists of other groups, without appropriate techniques for clarification required to obtain highly clarified slides.

Despite these problems, it was surprising that none of the studied species were new, suggesting that the Iranian knowledge of the group is better than expected. Of the 16 collected species, 12 were new records for the province, bringing the total of known species to 25. *Tydeus* was the most diverse genus with six species in our collections, including *Tydeus lindquisti* which was recorded for the first time from Iran. Although this species was first collected and described from litter under *Pinus* sp. in

Ontario, Canada (Marshall, 1970), it was later reported from bird nests from different parts of Poland (Kaźmierski *et al.* 2018). The two microscopic photographs (Fig.1, right and left) presented in this paper, are the only available photos of this species. The most abundant genera in the province were *Tydeus* with eight species and *Lorryia* with seven species. Future work on tydeoid mites will clarify the true nature of diversity and distribution of these mites: are tydeid mite species widespread, or is there an undetected endemic fauna in Mazandaran?

## REFERENCES

- Ahmad-Hosseini, M., Khanjani, M. & Karamian, R. (2019) Study on predatory mites of *Aceria tristriatus* (Nalepa, 1890) from Hamedan and Lorestan provinces, Western Iran. *Persian Journal of Acarology*, 8(2): 125–145.
- Akbari, A., Haddad Irani-Nejad, K., Khanjani, M., Arzanlou, M. & Kaźmierski, A. (2015) *Tydeus shabestariensis* sp. nov. and description of the male of *Neopronematus sepašgosariani* (Acari: Tydeoidea), with a key to the Iranian species of *Tydeus*. *Zootaxa*, 4032(3): 264–276.
- André, H.M. (1980) A generic revision of the family Tydeidae (Acari: Actinedida). IV. Generic description, key and conclusion. *Annales de la Societe Royale Zoologique de Belgique*, 116: 103–168.
- André, H.M. (1981) A generic revision of the family Tydeidae (Acari: Actinedida). III. Organotoaxy of the legs. *Acarologia*, 22: 165–178.
- André, H.M. & Fain, A. (2000) Phylogeny, ontogeny and adaptive radiation in the superfamily Tydeoidea (Acari: Actinedida), with a reappraisal of morphological characters. *Zoological Journal of the Linnean Society*, 130: 405–448.
- Baker, E.W. (1943) Nuevos Tydeidae Mexicanos (Acarina). *Revista de la Sociedad Mexicana de Historia Natural*, 4: 181–90.
- Baker, E.W. (1968) The genus *Paralorryia*. *Annals of Entomological Society of America*, 61: 1097–1106.
- Calvert, D.J. & Huffaker, C.B. (1974) Predator (*Metaseiulus occidentalis*) - prey (*Pronematus* spp.) interactions under sulfur and cattail pollen applications in a non-commercial vineyard. *Entomophaga*, 19: 361–369.
- Darbemamieh, M., Hajiqanbar, H., Khanjani, M. & Kaźmierski, A. (2015) New species and records of *Neopronematus* (Acari: Iolinidae) from Iran with a key to world species. *Zootaxa*, 3990(2): 235–246.
- Darbemamieh, M., Hajiqanbar, H., Khanjani, M. & Kaźmierski, A. (2016a) *Paurotyndareus*, a new genus of the family Iolinidae (Acari: Prostigmata), with the description of a new species from Iran. *Systematic and Applied Acarology*, 21(4): 398–404.
- Darbemamieh, M., Hajiqanbar, H., Khanjani, M., Gwiazdowicz, D. & Kaźmierski, A. (2016b) Some *Tydeus* mites (Acariformes: Prostigmata: Tydeidae) of Kermanshah province, western Iran, with remarks on *Tydeus caudatus*. *Acarologia*, 56(4): 603–611.
- Darbemamieh, M., Kamali, K. & Fathipour, Y. (2010) First report of *Tydeus caudatus* (Acari: Tydeidae) from Iran. *Journal of Entomological Society of Iran*, 30(1): 63–65.
- English-Loeb, G.M., Norton, A.P., Gadoury, D.M., Seem, R.C. & Wilcox, W.W. (1999) Control of powdery mildew in wild and cultivated grapes by a tydeid mite. *Biological Control*, 14: 97–103.
- Flaherty, D.L. & Hoy, M.A. (1971) Biological control of Pacific mites and Willamette mites in San Joaquin valley vineyards: Part III. Role of tydeid mites. *Researches in Population Ecology*, 13: 80–96.
- Fleschner, C.A. & Arakawa, K.Y. (1953) The mite *Tydeus californicus* on citrus and avocado leaves. *Journal of Economic Entomology*, 45(6): 1092.
- Kamali, K., Ostovan, H. & Atamehr, A. (2001) *A catalog of the mites and ticks (Acari) of Iran*. Islamic Azad Scientific Publication Centre, 192 pp.

- Kaźmierski, A. (1989) Revision of the genera *Tydeus* Koch sensu André, *Homeotydeus* André and *Orthotydeus* André with description of a new genus and four new species of Tydeinae (Acari: Actinedida: Tydeidae). *Mitteilungen aus dem Hamburgischen Zoologischen Museum und Institut*, 86: 289–314.
- Kaźmierski, A. (1998) Tydeinae of the world: generic relationships, new and redescribed taxa and keys to all species. A revision of the subfamilies Pretydeinae and Tydeinae (Acari: Actinedida: Tydeidae) - part IV. *Acta Zoologica Cracoviensia*, 41: 283–455.
- Kaźmierski, A. (2008) Tydeidae Kramer, 1877 sensu André et Fain, 2000 (Acari: Prostigmata). In: Bogdanowicz, W., Chudzicka, E., Pilipiuk, I. & Skibińska, E. (Eds.), *Fauna of Poland - Characteristics and checklist of species*. Zoological Museum and Institute of the Polish Academy of Science (PAN), Warsaw, 3, pp. 193–196 + 222 (In Polish).
- Kaźmierski, A. & Sikora, B. (2008) On the morphological anomalies in Tydeoidea (Actinedida). In: Bertrand, M., Kreiter, S., McCoy, K.D., Migeon, A., Navajas, M., Tixier, M.-S. & Vial, L. (Eds.), *Intergrative Acarology, Proceedings of the 6th European Congress, Montpellier, France*, pp. 82–88.
- Kaźmierski, A., Marciniak, M. & Sikora, B. (2018) Tydeinae mites (Acariformes: Prostigmata: Tydeidae) from bird nests with description of three new species. *Systematic and Applied Acarology*, 23(5): 803–823.
- Khanjani, M. & Ueckermann, E.A. (2003) Four new tydeid species from Iran (Acari: Prostigmata). *Zootaxa*, 182: 1–11.
- Marshall, V.G. (1970) Tydeid mites (Acarina: Prostigmata) from Canada. I. New and redescribed species of *Lorryia*. *Annals of Entomological Society of Quebec*, 15: 17–52.
- Mondin, S.A., Nuvoloni, F.M. & Feres, R.J.F. (2016) Four new species of *Lorryia* associated with *Hevea brasiliensis* Muell. Arg. (Euphorbiaceae) in Brasil. *Zootaxa*, 4158(4): 473–490.
- Niemczyk, E. & Kaźmierski, A. (2002) Checklist of species, occurrence and feeding of the members of families Stigmaeidae and Tydeidae in the apple orchards. In: Ignatowicz, S. (Ed.), *Postępy Polskiej Akarologii*, SGGW, Warszawa, pp. 265–274. (In Polish).
- Oudemans, A.C. (1928) Acarologische Aanteekeningen XCIV. *Entomologische Berichten*, 7: 374–382.
- Oudemans, A.C. (1929) Kritisch historisch Overzicht der Acarologie. Tweede Gedeelte. *Tijdschrifte voor Entomologie*, 72 (Suppl): I–XVII, 1–1097.
- Panou, H.N., Emmanouel, N.G. & Kaźmierski, A. (2000) *Neopronematus*, a new genus of the subfamily Pronematinae (Acari: Prostigmata: Tydeidae) and a new species from Greece. *Acarologia*, 41(3): 321–325.
- Sadeghi, H., Łaniecka, I. & Kaźmierski, A. (2012) Tydeoid mites (Acari: Triophtydeidae, Iolinidae, Tydeidae) of Razavi Khorasan Province, Iran, with description of three new species. *Annales Zoologici*, 62(1): 99–114.
- Sadeghi Namaghi, H. (1995) Survey of mites (Acari) fauna of pomaceous fruit trees in Mashhad region. *Agricultural Science and Technology Journal*, 9(1): 110–120.
- Silva, G.L., Metzelthin, M.H., Silva, O.S. & Ferla, N.J. (2016) Catalogue of the mite family Tydeidae (Acari: Prostigmata) with the world key to the species. *Zootaxa*, 4135(1): 1–68.
- Treat, A.E. (1970) Two tydeid mites from the ears of noctuid moths. *American Museum Novitates*, 2426: 1–14.
- Walter, D. & Krantz, G.W. (2009) Collection, rearing and preparing specimens. In: Krantz, G.W. & Walter, D.E. (Eds.), *A Manual of Acarology*. 3rd Edition. Texas Tech University, Lubbock, pp. 83–96.
- Zhang, Z.-Q., Fan, Q.-H., Pesic, V., Smith, H., Bochkov, A.V., Khaustov, A.A., Baker, A., Wohltmann, A., Wen, T., Amrine, J.W., Beron, P., Lin, J., Gabryś, G. & Husband, R. (2011)

Order Trombidiformes Reuter, 1909. *In*: Zhang, Z.-Q. (Ed.), *Animal Biodiversity: An outline of higher-level classification and survey of taxonomic richness*. *Zootaxa*, 3148: 129–138.

**COPYRIGHT**

Darbemamieh *et al.* Persian Journal of Acarology is under a free license. This open-access article is distributed under the terms of the Creative Commons-BY-NC-ND which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original author and source are credited.

## گزارش‌ها و نکته‌های جدیدی از کنه‌های بالاخانواده Tydeoidea (Trombidiformes: Prostigmata) از استان مازندران ایران

مریم درب‌امامیه<sup>۱\*</sup>، آندری کازمیرسکی<sup>۲</sup> و سعید پاک‌طینت سٹیج<sup>۳</sup>

۱- گروه گیاهپزشکی، دانشکده کشاورزی، پردیس کشاورزی و منابع طبیعی، دانشگاه رازی، کرمانشاه، ایران؛ رایانامه:

[m.darbemamieh@razi.ac.ir](mailto:m.darbemamieh@razi.ac.ir)

۲- گروه ریخت‌شناسی جانوری، دانشکده زیست‌شناسی، دانشگاه آدم میسکیویچ، پوزنان، لهستان؛ رایانامه: [akazmierski@tlen.pl](mailto:akazmierski@tlen.pl)

۳- گروه گیاهپزشکی، دانشکده علوم زراعی، دانشگاه علوم کشاورزی و منابع طبیعی ساری، ساری، ایران؛ رایانامه:

[saeedpaktinat@yahoo.com](mailto:saeedpaktinat@yahoo.com)

\* نویسنده مسئول

### چکیده

در این بررسی، ۱۶ گونه از شش جنس و دو خانواده از بالاخانواده Tydeoidea از استان مازندران معرفی می‌شوند. نمونه‌های مورد مطالعه، از دو کلکسیون متفاوت است، که توسط نویسنده نخست و سوم از استان مازندران جمع‌آوری شده‌اند. گزارشی جدید برای فون کنه‌های ایران، *Tydeus lindquisti* (Marshall, 1970)، دوازده گزارش جدید از استان مازندران و برخی نکته‌ها، توضیح‌ها و تصویرها در مورد این کنه‌ها آورده شده است.

واژگان کلیدی: *Tydeus lindquisti*، *Pronematus*، *Proctotydaeus*، *Neopronematus*، *Lorryia*، Iolinidae، *Homeopronematus*.

اطلاعات مقاله: تاریخ دریافت: ۱۳۹۸/۱۲/۱۹، تاریخ پذیرش: ۱۳۹۹/۲/۴، تاریخ چاپ: ۱۳۹۹/۴/۲۵