

## Article

### Phytoseiid mites (Acari: Phytoseiidae) of fruit orchards in cold regions of Razavi Khorasan province (northeast Iran), with redescription of two species

Hosnie Panahi Laeen<sup>1\*</sup>, Alireza Askarianzadeh<sup>1</sup> & Mahdi Jalaeian<sup>2</sup>

*1 Department of Plant Protection, Faculty of Agricultural Sciences, Shahed University, Tehran, Iran; E-mail: hosniepanahi@yahoo.com; askarianzadeh@shahed.ac.ir*

*2 Department of Plant Protection, Rice Research Institute of Iran (RRII), Rasht, Iran; E-mail: mahdi\_jalaeian@yahoo.com*

\*Corresponding Author

#### Abstract

Seven species from five genera of the family Phytoseiidae were collected in northeast Iran. *Typhlodromus (Anthoseius) neyshabouris* (Denmark & Daneshvar, 1982) were recorded for the second time. This species with the male of *Proprioiseiopsis messor* (Wainstein, 1960) are redescribed and illustrated. A key to the adult females of the Razavi Khorasan province of Iran is also provided. *Phytoseius corniger* Wainstein, 1959 had the highest abundance and distribution in this survey.

**Key words:** Predatory mite fauna, abundance, Mesostigmata, northeast Iran.

#### Introduction

Razavi Khorasan province is located in northeastern Iran. Mashhad is located in the center and is the capital of the province. Agriculture in Razavi Khorasan province is one of the largest and most important suppliers of agricultural products, with more than 1.06 million hectares under cultivation and horticultural crops play a decisive role in the economy of the province and country (Anonymous 2012). Predatory mites of the family Phytoseiidae are the most important natural enemies of tetranychid and eriophyid mites (Acari: Tetranychidae and Eriophyidae) (Gerson *et al.* 2003; Sabelis 1996). These mites feed on small insects such as whiteflies, thrips and scale insects as well as injurious plant mites. This family is relatively well known in Iran and about 70 species have been reported (Khalil-Manesh 1973; McMurtry 1977; Sepasgosarian 1977; Daneshvar 1980, 1987; Daneshvar & Denmark 1982; Kamali *et al.* 2001; Hajizadeh *et al.* 2002, 2009; Kolodochka *et al.* 2003; Shirdel 2003; Abbasipour *et al.* 2005; Rahmani *et al.* 2006, 2010; Faraji *et al.* 2007a, b, 2008a, b; Hajizadeh 2007; Shirdel *et al.* 2008; Ueckermann *et al.* 2009; Noei *et al.* 2010; Jafari *et al.* 2011; Asali Fayaz & Khanjani 2012, 2013; Asali Fayaz *et al.* 2012, 2013; Hajizadeh & Nazari 2012; Ostovan *et al.* 2012). In this study, a list of phytoseiid species collected from the fruit orchards (including apple, quincunx, pear and cherry) of the cold regions in the Razavi Khorasan province and a key to identification are presented.

## Materials and methods

This study was carried out in fruit orchards in Razavi Khorasan province, during 2009 and 2010. Phytoseiid mites were collected by beating or shaking shoots over a white tray or by extracting them from soil samples and fallen fruits using Berlese funnel. The mites were preserved in 70% ethanol and cleared in Nesbitt's fluid and then mounted on microscope slides using Hoyer's medium (Walter & Krantz 2009) for examination under an Olympus BX 41 phase contrast microscope. All specimens were collected by senior author. All measurements are given in micrometers ( $\mu\text{m}$ ). The classification systems used follows that of Chant & McMurtry (2007). The dorsal and ventral setal nomenclature is that of Rowell *et al.* (1978) and Chant & Yoshida-Shaul (1991). Idiosomal setal patterns are given according to Chant & Yoshida-Shaul (1992). All specimens are deposited in the Acarological Collection at Department of Plant Protection, Faculty of Agriculture, University of Shahed, Tehran, and in Jalal Afshar Zoological Museum, Department of Plant Protection, Faculty of Agriculture, University of Tehran, Karaj, Iran.

## Results

### Amblyseiinae Muma, 1961

#### *Proprioseiopsis messor* (Wainstein, 1960) (Figs. 1–8)

Syn.: *Typhlodromus messor* Wainstein, 1960: 668; *Amblyseius* (*Amblyseius*) *apheles* van der Merwe, 1968: 121 synonymy according to Ueckermann & Loots (1988); *Amblyseius lindquisti* Schuster & Pritchard, 1963: 246 synonymy according to Abbasova (1972).

#### *Distribution*

Algeria, Armenia, Australia, Azerbaijan, France, Gaza Strip, Greece, Israel, Italy, Morocco, New Zealand, South Africa, Spain, Turkmenistan, Ukraine and Iran (Asali Fayaz & Khanjani 2012; Ostovan *et al.* 2012), and new record for the region.

#### *Specimens examined*

Two adult females and one male from North Eastern Iran, Razavi Khorasan province, Torghabe–Shandiz, Kang village ( $36^{\circ} 19' 08'' \text{N}$ ,  $59^{\circ} 13' 36'' \text{E}$ ), 15 June 2009 ( $n=2$ ) and 21 May 2009 ( $n=1$ ). All from apple trees.


*Male* (Figs. 1–8). One specimen measured.

*Dorsum* (Fig. 2). Dorsal shield 312 long and 226 at the widest level, smooth with spots (not rugose patches) and pores on dorsal shield. Dorsal setae smooth, except for  $Z_5$ , serrate and setae  $J_5$  with two distinctive spines at base (Fig. 1); Setal lengths:  $j_1$  26,  $j_3$  53,  $j_4$  6,  $j_5$  5,  $j_6$  6,  $J_5$  10,  $z_2$  31,  $z_4$  21,  $z_5$  6,  $Z_1$  6,  $Z_4$  103,  $Z_5$  144,  $s_4$  85,  $S_2$  11,  $S_4$  7,  $S_5$  16;  $r_3$  20 and  $R_1$  8 on dorsal shield.

*Venter* (Fig. 3 & 5). Length of tritosternum (Fig. 5) 78, widened at its base 15. Sternal shield smooth and with two pairs of lyrifissures, length 145, width at widest point 125; with five pairs of setae:  $St_1$  31,  $St_2$  32,  $St_3$  28,  $St_4$  27,  $St_5$  29; ventrianal shield reticulated, with three pairs of pores, length 135, width at widest point 162; with 3 pairs of preanal setae  $JV_1$  25,  $JV_2$  18,  $ZV_2$  26;  $JV_5$  50, Para-anal setae 16 and Post-anal setae 17.

*Gnathosoma* (Fig. 4). Three pairs of smooth hypostomal setae, Length of setae:  $h_1$  20,  $h_2$  24,  $h_3$  22 and palp coxa with a pair of smooth setae,  $pc$  26. Corniculi 29 horn-like

and stout, internal malae somewhat longer than corniculi. Deutosternal groove with six transverse rows, basal row concave.



**Figures 1–3.** *Proprioseiopsis messor* (Wainstein, 1960) (male). 1. Dorsal setae J<sub>5</sub>; 2. Dorsal view of idiosoma; 3. Sternal and ventrianal shields.

*Palp.* Length of palp: 157, ratio of palp-tibia/tarsus length 1.39.

*Chelicera* (Fig. 6). Fixed digit 30 long with 3 teeth and a setiform pilus dentilis; movable digit 28 long with one tooth.


*Spermatodactyl.* L-shaped with a long foot. The outer margin of the shaft carries a membranous border.

*Peritreme* (Fig. 8). Extending anterior to setae j<sub>1</sub>, 185 long.

*Legs* (Fig. 7). Length of legs I–IV as follows: leg I 414, leg II 330, leg III 335 and leg IV 424. Leg IV with three macrosetae, on genu 67, tibia 52 and basitarsus 72. Genua I–IV with 10-8-7-7 setae.

#### Remarks

This redescription of the male is similar to the original description of Amitai & Wysoki (1974). However, it differs from their description in: dorsal setae Z<sub>4</sub> are smooth vs. serrated; dorsal setae J<sub>5</sub> with two distinctive spine at the base vs. simple; sternal shield with five pairs of setae (St<sub>1–5</sub>) vs. four pairs of setae (St<sub>1–4</sub>). Also, setae z<sub>5</sub>, s<sub>4</sub>, Z<sub>4</sub>, Z<sub>5</sub> and macrosetae on leg IV are longer than those of the Israeli specimens. Furthermore the Iranian specimens differ from specimens collected by Schicha (1983) as follows: shape of setae J<sub>5</sub> and setae j<sub>3</sub>, z<sub>2</sub>, z<sub>4</sub>, S<sub>2</sub> and macrosetae on leg IV of the Iranian specimens are longer than those of the Australian specimens.



**Figures 4–8.** *Proprioseiopsis messor* (Wainstein, 1960) (male). 4. Subcapitulum; 5. Tritosternum; 6. Chelicera & Spermatodactyl; 7. Leg IV; 8. Peritreme and peritremal plate.

### **Typhlodrominae Scheuten, 1857**

#### ***Typhlodromus (Anthoseius) neyshabouris* (Denmark & Daneshvar, 1982) (Figs. 9–16)**

##### *Distribution*

This species has been reported from Iran (Daneshvar & Denmark 1982) and here recorded for the second time from Iran and world.


##### *Specimens examined*

Three adult females from North Eastern Iran, Razavi Khorasan province: Torghabeh-Shandiz, Kang village (36° 19' 08" N, 59° 13' 36" E), 1 Sep. 2009; All from apple trees.

*Dorsum* (Fig. 9). Dorsal shield reticulated, with 3 pairs of large pores and 3 pairs of small pores. Measurements are as follows: Dorsal shield 365 (355–370) long and 180 (168–190) wide;  $j_1$  22 (20–25),  $j_3$  33 (30–36),  $j_4$  21 (17–22),  $j_5$  20 (18–22),  $j_6$  28 (22–30),  $J_2$  26 (22–33),  $J_5$  5 (4–5);  $z_2$  28 (27–31),  $z_3$  33 (29–36),  $z_4$  34 (30–40),  $z_5$  23 (21–25),  $Z_4$  48 (45–50),  $Z_5$  64 (57–71);  $s_4$  37 (33–41),  $s_6$  41 (39–42),  $S_2$  45 (41–50),  $S_4$  42 (36–46),  $S_5$  33 (29–35);  $r_3$  32 (32–34);  $R_1$  36 (33–40). Seta  $Z_5$  pointed apically.

*Venter* (Fig. 10). Length of tritosternum (Fig. 12) 94 (88–102), widened at its base 13 (13–14), sternal shield smooth and with two pairs of setae ( $St_1$  and  $St_2$ ) and two pairs of lyrifissures; setae  $St_3$  and  $St_4$  each set on separated platelets. Length of sternal setae as follows:  $St_1$  32 (31–33),  $St_2$  32 (32–33),  $St_3$  31 (30–31),  $St_4$  33 (32–33),  $St_5$  27 (25–28). Genital shield 134 (122–141) long and 75 (65–80) at widest level, spermatheca saccular,

calyx of spermatheca (Fig. 15) 21 (18–24) long, with C-shaped atrium. Ventrianal shield 113 (106–127) long, 72 (68–76) at widest level; with a pair of pores and four of pairs preanal setae, setae JV<sub>1</sub> 18 (16–20), JV<sub>2</sub> 20 (18–22), JV<sub>3</sub> 21 (17–22), JV<sub>4</sub> 20 (20–21), JV<sub>5</sub> 55 (52–65), ZV<sub>1</sub> 23 (21–26), ZV<sub>2</sub> 20 (18–21), ZV<sub>3</sub> 14 (13–16), both para-anal and post-anal setae 16.



**Figures 9–10.** *Typhlodromus (A.) neyshabouris* (Denmark & Daneshvar, 1982) (female). 9. Dorsal view of idiosoma; 10. Ventral view of idiosoma.

*Gnathosoma* (Fig. 11). With three pairs of smooth hypostomal setae. Lengths of setae: h<sub>1</sub> 23 (19–25), h<sub>2</sub> 20 (16–23), h<sub>3</sub> 21 (20–23) and palp coxa with a pair of smooth capitular setae, pc 30 (28–32). Corniculi 23 (22–24), horn-like and stout, corniculi somewhat longer than internal malae.

*Palp.* Length of palp: 183 (181–185), ratio of palp-tibia/tarsus length 1.41 (1.3–1.46).

*Chelicera* (Fig. 14). Movable digit 27 (26–28), with a tooth; fixed digit 30 (28–31), with four teeth and a *pilus dentilis*.

*Peritreme* (Fig. 13). 49 (44–55) long. Apex of peritreme extending close to setae s<sub>4</sub> and s<sub>6</sub>, but not to z<sub>2</sub>.


*Legs* (Fig. 16). Length of legs I–IV as follows: leg I 314 (305–320), leg II 274 (265–289), leg III 293 (282–295) and leg IV 376 (371–384). Basitarsus IV with one macroseta 57 (52–61). Genua I–IV with 10-7-7-7 setae.

*Male:* Unknown.

#### Remarks

The characteristics of the specimens collected are very similar to those of the original

description of Denmark & Daneshvar (1982). However, it differs from them in: fixed digit of chelicera with four teeth plus a *pilus dentilis* vs. three teeth plus a *pilus dentilis* in the original description. Also, length and width of dorsal shield and all setae of these specimens are longer than those of the original description.



**Figures 11–16.** *Typhlodromus (A.) neyshabouris* (Denmark & Daneshvar, 1982). 11. Gnathosoma; 12. Tritosternum; 13. Peritreme; 14. Chelicera; 15. Spermatheca; 16. Leg IV.

***Typhlodromus (A.) bagdasarjani* (Wainstein & Artunjan, 1967)**

*Typhlodromus bagdasarjani* Wainstein & Arutunjan, 1967: 1765; de Moraes *et al.* 2004: 311; *Typhlodromus (Anthoseius) kettanehi* Dosse, 1967: 32.

*Distribution*

Armenia, Azerbaijan, Iran, Lebanon, Turkey and Turkmenistan (Rahmani *et al.* 2010).

*Specimens examined*

Razavi Khorasan province: Laeen village, Kalat Naderi (37° 07' 45" N, 59° 29' 56" E), 19 Oct. 2009 (n= 1); Kang village, Torghabe–Shandiz (36° 19' 08" N, 59° 13' 36" E), 2 Sep. 2009, (n= 5); Torogh, Mashhad (36° 12' 37" N, 59° 39' 36" E), 2 Sep. 2009 (n= 3) and 16 Oct. 2009 (n= 2); Mashhad (36° 18' 28" N, 59° 31' 36" E), 15 Sep. 2009 (n= 22) and 13 Oct. 2009 (n= 5); All collected on apple, peach, quince and cherry trees.

**Phytoseiinae Berlese, 1916**

***Phytoseius corniger* Wainstein, 1959**

*Distribution*

Azarbaijan, Iran, Turkey and Turkmenistan (Hajizadeh 2007).

### *Specimens examined*

Razavi Khorasan province: Laeen village, Kalat Naderi (37° 07' 45" N, 59° 29' 56" E), 13 Oct. 2009 (n= 56), 5 Nov. 2009 (n= 28), 2 Sep. 2009 (n= 49); Kang village, Torghabe–Shandiz (36° 19' 08" N, 59° 13' 36" E), 11 Sep. 2009 (n= 4); Torogh, Mashhad (36° 12' 37" N, 59° 39' 36" E), 1 Oct. 2009 (n= 35) 15 Sep. 2009 (n= 19); Mashhad, (36° 18' 28" N, 59° 31' 36" E), 24 May 2009 (n= 6), 20 June 2009 (n= 4), 22 Aug. 2009 (n= 10), 1 Sep. 2009 (n= 75), 8 Sep. 2009 (n= 58), 15 Sep. 2009 (n= 47), 19 Sep. 2009 (n= 142), 27 Sep. 2009 (n= 123), 7 Oct. 2009 (n= 102), 13 Oct. 2009 (n= 71), 25 Oct. 2009 (n= 96), 31 Oct. 2009 (n= 82), 6 Nov. 2009 (n= 45), 14 Nov. 2009 (n= 18), 12 Dec. 2009 (n= 12), 30 Apr. 2010 (n= 9), 6 May 2010 (n= 14); All on apple, peach and cherry trees.

### **Amblyseiinae Muma, 1961**

#### ***Neoseiulus barkeri* Hughes, 1948**

*Neoseiulus barkeri* Hughes, 1948: 141; *Amblyseius mckenziei* Schuster & Pritchard, 1963: 268; *Amblyseius (Amblyseius) usitatus* (van der Merwe, 1965): 71; *Amblyseius oahuensis* Prasad, 1968: 1518; *Amblyseius picketti* Specht, 1968: 681; *Amblyseius mycophilus* Karg, 1970: 290; *Amblyseius masiaka* Blommers & Chazeau, 1974: 308; *Neoseiulus kermanicus* Daneshvar, 1987: 14; Faraji *et al.* 2007a: 233.

#### *Distribution*

This species is known from Algeria, Australia, Brazil, China, England, Finland, France, Germany, Ghana, Greece, Guinea, Hawaii, India, Iran, Italy, Japan, Jordan, Madagascar, Mozambique, Nigeria, Palestine, Russia, Spain, South Africa, South Korea, Sweden, Taiwan, The Netherlands, Turkey, Ukraine, USA, Yemen (Rahmani *et al.* 2010), Canary Islands, Cape Verde, Georgia, Norway, Reunion Island and West Bank (Asali Fayaz & Khanjani 2012) and this is a new record for this region.

### *Specimens examined*

Razavi Khorasan province: Torghabe–Shandiz (36° 18' 37" N, 59° 22' 25" E), 6 Aug. 2009 (n= 1); on apple trees.

#### ***Euseius amissibilis* Meshkov, 1991**

*Euseius amissibilis* Meshkov 1991: 138.

#### *Distribution*

This species has been reported from Azarbaijan, Iran, Turkey, Turkmenistan (Hajizadeh 2007) and Tajikistan (Rahmani *et al.* 2010) and this is also a new record for this region.

### *Specimens examined*

Razavi Khorasan province: Laeen village, Kalat Naderi (37° 07' 45" N, 59° 29' 56" E), 31 Aug. 2009 (n= 1); 19 Oct. 2009 (n= 44), 29 Oct. 2009 (n= 95), 2 Nov. 2009 (n= 55), 12 Nov. 2009, (n= 64), 20 Nov. 2009 (n= 67); all on apple, peach and cherry trees.

#### ***Neoseiulus marginatus* (Wainstein, 1961)**

*Typhlodromus marginatus* Wainstein, 1961: 158; *N. polyporus* (Wainstein, 1962): 143.

### Distribution

This species has been reported from Algeria, Armenia, Azerbaijan, Georgia, Greece, France, Moldova, Russia, Turkmenistan, Ukraine (de Moraes *et al.* 2004), Hungary, Kazakhstan and Kenya (Asali Fayaz & Khanjani 2012) and a new record for this region.

### Specimens examined

Razavi Khorasan province: Mashhad (36° 18' 28" N, 59° 31' 36" E), 14 Nov. 2009 (n=1) and 10 Oct. 2009 (n=2); all on apple trees.


### Key to the Phytoseiidae of Razavi Khorasan province (northeast Iran) (female)

1. Seta  $z_3$  and  $s_6$  absent (Amblyseiinae Muma) .....6  
– Either or both setae  $z_3$  and  $s_6$  present .....2
2. Setae  $Z_1$ ,  $S_2$ ,  $S_4$  and  $S_5$  absent (Phytoseiinae Berlese, *Phytoseius* Ribaga) .....*P. corniger* Wainstein  
– At least one of setae  $Z_1$ ,  $S_2$ ,  $S_4$  and  $S_5$  present (Typhlodrominae Chant & McMurtry) .....3
3. Ventrianal shield with three pairs of preanal setae .....  
.....*Typhlodromus (Anthoseius) rodriguizi* (Denmark & Daneshvar)  
– Ventrianal shield with four pairs of preanal setae .....4
4. Dorsal shield with 5 pairs of large pores; movable digit of chelicerae without teeth .....*T. bagdasarjani* Wainstein & Arutunjan  
– Dorsal shield with 3 pairs of large pores; movable digit of chelicerae with one tooth .....5
5. Ventrianal shield with a pair of pores.....*T. (A.) neyshabouris* (Denmark & Daneshvar)  
– Ventrianal shield without any pores.....*T. (A.) torbatejamae* (Denmark & Daneshvar)
6. Seta  $JV_1$  inserted well behind anterior margin of ventrianal shield and preanal setae arranged in an almost transverse row; cheliceral digits short and stout (*Euseius* Wainstein) .....  
.....*Euseius amissibilis* Meshkov  
– Seta  $JV_1$  inserted near margin of ventrianal shield and preanal setae not arranged in a transverse row across the shield, cheliceral digits elongate.....7
7. Macrosetae present only on leg IV or absent and  $J_2$  present (*Neoseiulus* Hughes) .....8  
– Macrosetae present at least on genua III, as well as on leg IV and  $J_2$  absent.....  
.....*Proprioseiopsis messor* (Wainstein)
8. Spermatheca with a stalk between atrium and calyx.....*N. marginatus* (Wainstein)  
– Spermatheca without a stalk between atrium and calyx.....*N. bakeri* Hughes

### Discussion

In the study, seven species from five genera of the family Phytoseiidae were reported from Razavi Khorasan province (northeast Iran). *Typhlodromus (Anthoseius) neyshabouris* (Denmark & Daneshvar, 1982) were recorded for the second time in Iran and the male of *Proprioseiopsis messor* (Wainstein, 1960) are redescribed. *Phytoseius corniger* Wainstein, 1959 was the most abundant species among the seven species collected in fruit orchards in the cold regions of Razavi Khorasan province (northeast Iran) (Fig. 17). The species was collected from all sampled regions and has a good distribution in the studied area. Therefore, it is important to protect or conserve the

predator in the environment especially by reducing pesticide application in pest management program.



**Figure 17.** Relative abundance of phytoseiid predatory mites in fruit orchards of the cold regions of Razavi Khorasan province (northeast Iran) during surveys in 2009 and 2010.

### Acknowledgements

The authors wish to express their gratitude to Prof. Dr. Alireza Saboori (Department of Plant Protection, Faculty of Agriculture, University of Tehran, Karaj, Iran) for his advice, guidance, interest and motivation, and Dr. Farid Faraji (MITOX Consultants, Amsterdam, The Netherlands) for his kind help in the identification of the specimens. This project is a part of the M. Sc. thesis of the senior author and was financially supported by the Shahed University, which is greatly appreciated.

### References

- Abbasipour, H., Taghavi, A. & Ueckermann, E.A. (2005) Redescription of *Transeius patellae* (Karg) (Acari: Phytoseiidae) and first record from Iran. *International Journal of Acarology*, 31(4): 363–366.
- Abbasova, E.D. (1972) Phytoseiid mites (Parasitiformes: Phytoseiidae) of Azerbaijan. *Avtoreferat Dissertatsii na Soiskanie Uchenoy Stepeni Kandidata Biologicheskikh Nauk. Akademiya Nauk Azerbaydzhanskoy SSR, Institut Zoologii, Baku, Azerbaijan*, 34 pp.
- Amitai, S. & Wysoki, M. (1974) Two unknown males of genus *Amblyseius* Berlese and their karyotypes (Mesostigmata: Phytoseiidae). *Acarologia*, 16: 45–51.
- Anonymous, (2012) Agriculture in Razavi Khorasan Province. Jihad Agriculture of Razavi Khorasan Province Press, 22 pp. (In Persian).

- Asali Fayaz, B. & Khanjani, M. (2012) Phytoseiid mites (Acari: Mesostigmata: Phytoseiidae) in some regions of western and north western Iran. *Journal of Crop Protection*, 1(2): 161–172.
- Asali Fayaz, B. & Khanjani, M. (2013) Redescription of four species of phytoseiid mites (Acari: Mesostigmata) associated with alfalfa farms in western Iran. *Persian Journal of Acarology*, 2(1): 9–24.
- Asali Fayaz, B., Khanjani, M., Hajizadeh, J. & Ueckermann, E.A. (2012) A re-description of *Typhlodromus (Anthosieus) Tamaricis* (Kolodochka) (Mesostigmata: Phytoseiidae), a first record for Iran. *Acarologia*, 52(4): 425–431.
- Asali Fayaz, B., Khanjani, M. & Tixier M. (2013) Redescription of six species of the genus *Typhlodromus* Scheuten (Acari: Phytoseiidae: Typhlodrominae) recorded from some regions of Western and North-Western Iran. *Persian Journal of Acarology*, 2(3): 369–387.
- Blommers, L. & Chazeau, J. (1974) Two new species of predator mites of the genus *Amblyseius* Berlese (Acarina: Phytoseiidae) from Madagascar. *Zeitschrift für Angewandte Entomologie*, 75: 308–315.
- Chant, D.A. & McMurtry, J.A. (2007) Illustrated keys and diagnoses for the genera and subgenera of the Phytoseiidae of the world (Acari: Mesostigmata). *Indira Publishing House, West Bloomfield*, 220 pp.
- Chant, D.A. & Yoshida-Shaul, E. (1991) Adult ventral setal patterns in the family Phytoseiidae (Acari: Gamasina). *International Journal of Acarology*, 17 (3):177–193.
- Chant, D.A. & Yoshida-Shaul, E. (1992) Adult idiosomal setal patterns in the family Phytoseiidae (Acari: Gamasina). *International Journal of Acarology*, 18 (3):187–199.
- Daneshvar, H. (1980) Some predator mites from northern and western Iran. *Applied Entomology and Phytopathology*, 48: 15–17.
- Daneshvar, H. (1987) Some predatory mites from Iran, with descriptions of one new genus and six new species (Acari: Phytoseiidae, Ascidae). *Applied Entomology and Phytopathology*, 54 (1–2): 13–37.
- Daneshvar, H. & Denmark, H.A. (1982) Phytoseiids of Iran (Acarina: Phytoseiidae). *International Journal of Acarology*, 8: 3–14.
- de Moraes, G.J., McMurtry, J.A., Denmark, H.A. & Campos, C.B. (2004) A revised catalog of the mite family Phytoseiidae. *Zootaxa*, 434: 494 pp.
- Dosse, G. (1967) Schadmilben des Libanons und ihre Pradatoren. *Zeitschrift für Angewandte Entomologie*, 59: 16–48.
- Faraji, F., Hajizadeh, J., Ueckermann, E.A., Kamali, K. & McMurtry, J.A. (2007) Two new records for Iranian phytoseiid mites with synonymy and keys to the species of *Typhloseiulus* Chant & MacMurtry and Phytoseiidae in Iran (Acari: Mesostigmata). *International Journal of Acarology*, 33 (3): 231–239.
- Faraji, F., Jalaeian, M. & McMurtry, J.A. (2008) A new species of *Paraseiulus* Muma from Iran with a key to the known species (Acari: Mesostigmata: Phytoseiidae). *Zootaxa*, 1770: 65–68.
- Faraji, F., Sakenin-Chelav, H., Kamali, K. & McMurtry, J.A. (2008) Four new species records of Phytoseiidae (Acari: Mesostigmata) for Iran, and description of variability in the spermatheca of *Typhlodromus bakeri*. *Systematic & Applied Acarology*, 13: 123–132.


- Faraji, F., Shiroodbakhshi, M., Ostovan, H. & McMurtry J.A. (2007) Redescription of the female of *Paraseiulus triporus* and *Proprioseiopsis dacus* (Acari: Phytoseiidae) based on material collected from citrus in northern Iran. *Systematic & Applied Acarology*, 12: 199–204.
- Gerson, U., Smiley, R.L. & Ochoa, R. (2003) *Mites (Acari) for Pest Control*. Blackwell Science Ltd., UK. 539 pp.
- Hajizadeh, J. (2007) Introducing a part of the phytoseiids (Acari: Phytoseiidae) fauna of Guilan province, part II: subfamily Amblyseiinae Muma and Phytoseiinae Berlese. *Agriculture Research*, 7 (1): 7–25 (in Persian with English abstract).
- Hajizadeh, J., Faraji, F. & Rafatifard, M. (2009) *Predatory mites of the family Phytoseiidae of Iran*. University of Guilan Press, 282 pp.
- Hajizadeh, J., Hosseini, R. & McMurtry, J. A. (2002) Phytoseiid mites (Acari: Phytoseiidae) associated with eriophyid mites (Acari: Eriophyidae) in Guilan province of Iran. *International Journal of Acarology*, 28 (4): 373–378.
- Hajizadeh, J. & Nazari, M. (2012) A checklist and key for the phytoseiid mites (Acari: Phytoseiidae) of citrus orchards in Iran, with a new record for Iranian phytoseiid mites. *Systematic & Applied Acarology*, 17 (4): 388–396.
- Hughes, A.M. (1948) *The mites associated with stored food products*. Ministry of Agriculture and Fisheries, H. M. Stationary Office, London, 168 pp.
- Jafari, Sh., Fathipour, Y. & Faraji, F. (2011) Re-descriptions of *Amblyseius meghriensis* Arutunjan and *Typhlodromus haiastanius* (Arutunjan) with discussion on using preanal pores as a character in the subgenus *Anthoseius* (Mesostigmata: Phytoseiidae). *International Journal of Acarology*, 37 (3): 244–254.
- Kamali, K., Ostovan, H. & Atamehr, A. (2001) *A catalog of mites & ticks (Acari) of Iran*. Islamic Azad University Scientific Publication Center, 206 pp.
- Karg, W. (1970) Neue Arten der Raubmilbenfamilie Phytoseiidae Berlese, 1916 (Acarina: Parasitiformes). *Deutsche Entomologische Zeitschrift*, 17: 289–301.
- Khalil-Manesh, B. (1973) Phytophagous mites fauna of Iran. *Applied Entomology and Phytopathology*, 35: 30–38 (in Persian with English abstract).
- Kolodochka, L.A., Hajiqanbar, H. & McMurtry, J.A. (2003) A description of unknown male and redescription of female of the rare Phytoseiid mite *Neoseiulus sugonjaevi* (Wainstein & Abbasova, 1974) (Parasitiformes, Phytoseiidae) from Iran. *Acarina*, 11(2): 231–233.
- McMurtry, J.A. (1977) Description and biology of *Typhlodromus persianus* n. sp., from Iran with notes on *T. kettanehi* (Acarina: Mesostigmata: Phytoseiidae). *Annals of the Entomological Society of America*, 70: 563–568.
- Meshkov, Y.I. (1991) *Euseius amissibilis* sp. n. (Parasitiformes, Phytoseiidae), a new species of mite from Tadjikistan [in Russian]. *Zoologicheskii Zhurnal*, 70: 138–140.
- Noei, J., Hajizadeh, J., Faraji, F., Ostovan, H. & Salehi, L. (2010) First report of two species of predatory mites (Acari: Phytoseiidae) for Iranian fauna. *Journal of Entomological Society of Iran*, 29 (2): 113–116.
- Ostovan, H., Faraji, F., Kamyab, F. & Khadempour, F. (2012) Notes on *Neoseiulus paspalivorus* (Deleon) and *Proprioseiopsis messor* (Wainstein) (Acari: Phytoseiidae) collected in Iran. *Acarologia*, 52 (1): 51–58.
- Prasad, V. (1968) *Amblyseius* mites from Hawaii. *Annals of the Entomological Society of America*, 61 (6): 1514–1521.
- Rahmani, H., Kamali, K. & Faraji F. (2010) Predatory mite fauna of Phytoseiidae of northwest Iran (Acari: Mesostigma). *Turkish Journal of Zoology*, 34: 497–508.

- Rahmani, S., Saboori, A., Ueckermann, E. & Ardeshir, F. (2006) Fauna of phytoseiid mites (Mesostigmata) in orchards of Karaj region, Iran. *Abstract book of 12<sup>th</sup> International Congress of Acarology, Amsterdam, The Netherlands*, p. 167.
- Rowell, H.J., Chant, D.A. & Hansell, R.I.C. (1978) The determination of setal homologies and setal patterns on the dorsal shield in the family Phytoseiidae (Acarina: Mesostigmata). *Canadian Journal of Entomology*, 110: 859–876.
- Sabelis, M. W. (1996) *Phytoseiidae*. In: Lindquist, E. E., Sabelis, M. W. & Bruin, J. (Eds.) *Eriophyoid mites, their biology, natural enemies and control*. Elsevier Science Publishing, Amsterdam, The Netherlands, pp. 427–456.
- Schicha, E., (1983) New species, new records, and redescription of phytoseiid mites from Australia, Tahiti and the African region (Acari: Phytoseiidae). *International Journal of Entomology*, 25(2–3): 103–126.
- Sepasgosarian, H. (1977) The 20 years research of Acarology in Iran. *Journal of the Iranian Society of Engineers*, 56: 40–50.
- Shirdel, D. (2003) *Survey on the phytoseiid fauna of East Azarbaijan*. Ph. D. thesis. Department of Entomology, Science and Research Campus, Islamic Azad University, Tehran, Iran, 183 pp.
- Shirdel, D. Kamali, K. & Faraji, F. (2008) Redescription of *Typhloseiulus carmonae* (Chant and Yoshida-Shaul) (Mesostigmata: Phytoseiidae) new species for Iran. *Acarina*, 16 (1):51–56.
- Schuster, R.O. & Pritchard, A.E. (1963) Phytoseiid mites of California. *Hilgardia*, 34: 191–285.
- Specht, H.B. (1968) Phytoseiidae (Acarina: Mesostigmata) in the New Jersey apple orchard environment with descriptions of spermathecae and three new species. *Canadian Entomologist*, 100: 673–692.
- Ueckermann, E.A. & Loots, G.C. (1988) *The African species of the subgenera Anthoseius De Leon and Amblyseius Berlese (Acari: Phytoseiidae)*. Entomology Memoir, Department of Agriculture and Water Supply, Republic of South Africa, No. 73, 168 pp.
- Ueckermann, E.A., Jalaieian, M., Saboori, A. & Seyedoleslami, H. (2009) Re-description of *Typhlodromus (Anthoseius) khosrovensis*, first record for Iran (Acari: Phytoseiidae). *Acarologia*, 49: (1–2): 23–27.
- van der Merwe, G.G. (1965) South African Phytoseiidae (Acarina). I. Nine new species of the genus *Amblyseius* Berlese. *Journal of the Entomological Society of South Africa*, 28: 57–76.
- van der Merwe, G.G. (1968) *A taxonomic study of the family Phytoseiidae (Acari) in South Africa with contributions to the biology of two species*. Entomology Memoirs, South Africa Department of Agricultural Technical Services, South Africa, 18: 1–198.
- Wainstein, B.A. (1959) New subgenus and species of the genus *Phytoseius* Ribaga, 1902 (Phytoseiidae: Parasitiformes). *Zoologicheskii Zhurnal*, 38: 1361–1365.
- Wainstein, B.A. (1960) New species and subspecies of the genus *Typhlodromus* Scheuten (Parasitiformes, Phytoseiidae) of the USSR fauna. *Zoologicheskii Zhurnal*, 39: 683–690.
- Wainstein, B.A. (1961) New species of mites of the genus *Typhlodromus* (Parasitiformes: Phytoseiidae) in Georgia. [in Russian]. *Trudy Instituta Zoologii Akademii Nauk Gruzinskoy SSR*, 18: 153–162.

- Wainstein, B.A. (1962) Some new predatory mites of the family Phytoseiidae (Parasitiformes) of the USSR fauna. *Annual Review of Entomology*, 41: 139–146 [English translation].
- Wainstein, B.A. & Arutunjan, E.S. (1967) New species of predaceous mites of the genera *Typhlodromus* Scheuten and *Paraseiulus* Muma (Parasitiformes, Phytoseiidae) [in Russian] *Zoologicheskii Zhurnal*, 46: 1764–1770.
- Walter, D.E. & Krantz, G.W. (2009) Collecting, rearing, and preparing specimens. In: Krantz, G.W. & Walter, D.E. (Eds.) *A manual of Acarology*, 3<sup>rd</sup> edition. Texas Tech University Press, pp. 83–96.

Received: 11 October 2013  
Accepted: 18 November 2013  
Published: 15 January 2014

#### COPYRIGHT

 Panahi *et al.* Persian Journal of Acarology is under free license. This open-access article is distributed under the terms of the Creative Commons-BY-NC-ND which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original author and source are credited.

### کنه‌های فیتوزئید (**Acari: Phytoseiidae**) باغ‌های میوه سردسیری استان خراسان رضوی (شمال شرق ایران) همراه با باز توصیف دو گونه

حسینیه پناهی لائین<sup>۱</sup>، علیرضا عسکریان‌زاده<sup>۱</sup> و مهدی جلائیان<sup>۲</sup>

<sup>۱</sup> گروه گیاهپزشکی، دانشکده علوم کشاورزی، دانشگاه شاهد، تهران، ایران؛ رایانامه: [hosniepanahi@askarianzadeh@shahed.ac.ir](mailto:hosniepanahi@askarianzadeh@shahed.ac.ir) yahoo.com

<sup>۲</sup> بخش گیاهپزشکی، موسسه تحقیقات برنج کشور، رشت، ایران؛ رایانامه: [mahdi\\_jalaeian@yahoo.com](mailto:mahdi_jalaeian@yahoo.com)

#### چکیده

هفت گونه از پنج جنس از خانواده Phytoseiidae از شمال شرق ایران جمع‌آوری شد. گونه *Typhlodromus (Anthoseius) neyshabouris* (Denmark & Daneshvar, 1982) برای دومین بار

گزارش می‌شود. این گونه به همراه نمونه نر گونه *Proprioseiopsis messor* (Wainstein, 1960) بازتوصیف و ترسیم شدند. کلیدی برای کنه‌های ماده استان خراسان رضوی ایران نیز تهیه شده است. گونه *Phytoseius corniger* Wainstein, 1959 بیشترین فراوانی و پراکندگی را در این پژوهش داشت.

**واژگان کلیدی:** فون کنه‌های شکارگر، فراوانی، میان‌استیگمایان، شمال شرق ایران.

تاریخ دریافت: ۱۳۹۲/۷/۱۹

تاریخ پذیرش: ۱۳۹۲/۸/۲۷

تاریخ چاپ: ۱۳۹۲/۱۰/۲۵