



First record of the genus *Phauloppia* (Acari, Oribatida, Oribatulidae) in Ethiopia, with description of a new arboreal species

Sergey G. Ermilov^{1*} | Vladimir M. Salavatulin² | Leonid B. Rybalov²

1. University of Tyumen, X-BIO Institute, Tyumen, Russia; E-mail: ermilovacari@yandex.ru

2. Severtsov Institute of Ecology and Evolution, Russian Academy of Sciences, Laboratory of Soil Zoology and General Entomology, Moscow, Russia; E-mails: v.salavatulin@gmail.com, rybalov52@mail.ru

* Correspondence

ermilovacari@yandex.ru

Received:

24 April, 2026

Accepted:

09 June, 2026

Published:

15 July, 2026

Subject Editor:

O. Jobarchi



<http://zoobank.org/urn:lsid:zoobank.org:pub:F6B55C07-40FE-42F5-892C-6B5D35CEF7AA>

ABSTRACT

The genus *Phauloppia* (Oribatida, Oribatulidae) is recorded for the first time from Ethiopia. A new species, *P. hageniaica* **sp. nov.**, is described based on females collected from bark and moss of *Hagenia abyssinica* in southeastern Ethiopia. The new species is characterized by the absence of the lamella.

KEYWORDS

Afrotropical region, morphology, oribatid mites, taxonomy, tree dweller

CITE: Ermilov, S.G., Salavatulin, V.M. & Rybalov, L.B. (2026) First record of the genus *Phauloppia* (Acari, Oribatida, Oribatulidae) in Ethiopia, with description of a new arboreal species. *Persian Journal of Acarology*, 15(3): 150310.
<https://doi.org/10.22073/pja.150310>

INTRODUCTION

The oribatid mite genus *Phauloppia* (Acari, Oribatida, Oribatulidae) comprises 25 species having a cosmopolitan distribution collectively (Subías 2022, online version 2024; Ermilov *et al.* 2024). Species of the genus inhabit various substrates across different ecosystems; some species are arboreal (e.g., Bayartogtokh 2010; Murvanidze and Mumladze 2016; Ermilov and Salavatulin 2025). Generic diagnosis was revised by Ermilov and Salavatulin (2023). Identification keys to some species of *Phauloppia* were presented by, e.g., Balogh and Balogh (2002), Weigmann (2006), Bayartogtokh (2010), and Ermilov and Salavatulin (2023).

The oribatid mite fauna of Ethiopia comprises 375 species; representatives of *Phauloppia* have not been recorded (Ermilov and Rybalov 2026). During taxonomic identification of arboreal mites from *Hagenia abyssinica* (Rosaceae) in southeastern Ethiopia, we found a new species belonging to *Phauloppia*. The main goal of the paper is to describe this new species under the name *Phauloppia hageniaica* **sp. nov.**

Eight species of Oribatulidae (1 species – *Lunoribatula* Mahunka, 1982(a); 2 species – *Oribatula* Berlese, 1896; 5 species – *Zygoribatula* Berlese, 1916) have been previously recorded in the Ethiopian fauna (Ermilov and Rybalov 2026).

METHODS

For measurement and illustration, specimens were mounted in lactic acid on temporary cavity



slides. Body length was measured in lateral view, from the tip of the rostrum to the posterior edge of the gastronotum. Maximum body width refers to the maximum width of the notogaster in dorsal view. Setae were measured from the side to avoid foreshortening. All measurements are in micrometres (μm). Formulas for leg setation are given in parentheses according to the sequence trochanter–femur–genu–tibia–tarsus (femulus of tarsus I included). Formulas for leg solenidia are given in square brackets, according to the sequence genu–tibia–tarsus. Paired structures are described in the singular, unless otherwise noted. Drawings were made with a camera lucida using a Leica DM 2500 light microscope.

Morphological terminology used in this paper follows that of Grandjean: see Trávě and Vachon (1975) for references, Norton (1977) for leg setal nomenclature, and Norton and Behan-Pelletier (2009) for overview.

Abbreviations: Prodorsum: *ro*, *le*, *in*, *bs*, *ex* = rostral, lamellar, interlamellar, bothridial, and exobothridial setae, respectively; *Ad* = dorsosejugal porose area; *D* = dorsophragma; *P* = pleurophragma. Notogaster: *c*₁, *c*₂, *da*, *dm*, *dp*, *la*, *lm*, *lp*, *b*₁–*b*₃, *p*₁–*p*₃ = setae; *Aa*, *A1*, *A2*, *A3* = porose areas; *ia*, *im*, *ip*, *ib*, *ips* = lyrifissures; *gla* = opisthonotal gland opening. Gnathosoma: *a*, *m*, *b* = subcapitular setae; *or* = adoral seta; *d*, *l*, *cm*, *acm*, *ul*, *su*, *lt*, *vt*, *inf*, *sup* = palp setae; ω = palp solenidium; *cha*, *chb* = cheliceral setae; *Tg* = Trägårdh's organ. Epimeral and lateral podosomal regions: *1a*, *1b*, *1c*, *2a*, *3a*, *3b*, *3c*, *4a*, *4b*, *4c* = epimeral setae; *PdI*, *PdII* = pedotecta I and II, respectively; *dis* = discidium; *cir* = circumpedial carina. Anogenital region: *g*, *ag*, *an*, *ad* = genital, aggenital, anal, and adanal setae, respectively; *iad* = adanal lyrifissure; *Ap* = postanal porose area; *po* = preanal organ. Legs: *pa* = porose area; *d*, *l*, *v*, *ev*, *bv*, *ft*, *tc*, *it*, *p*, *u*, *a*, *s*, *pv*, *pl* = setae; *e* = famulus; ω , σ , φ = solenidia.

RESULTS

TAXONOMY

Family Oribatulidae Thor, 1929

Genus *Phauloppia* Berlese, 1908

Type species: *Zetes lucorum* C.L. Koch, 1841 (syn.: *Oppia conformis* Berlese, 1895)

Phauloppia hageniaica sp. nov. (Figs 1–10)

<http://zoobank.org/urn:lsid:zoobank.org:act:6FF1A2F1-7444-40C2-B39E-0207F3634FC1>

Type material

Holotype (female) and two paratypes (two females): southeastern Ethiopia, Oromia Region, Bale Zone, Bale Mountains National Park, 6° 44' 37" N, 39° 43' 00" E, 2628 m a.s.l., Harennna Forest, forest with *Hagenia abyssinica* Willd. and *Oldeania alpina* (K. Schum.), bark and moss on *H. abyssinica* at a height of 7 m, 16.XII.2025 (collected by V.M. Salavatulin). Detailed description of arboreal acarofauna extraction techniques is presented in Salavatulin (2019).

Type deposition

The holotype is deposited in the collection of the Senckenberg Museum of Natural History, Görlitz, Germany. Two paratypes are deposited in the collection of the University of Tyumen, Museum of Zoology, Tyumen, Russia. All specimens are preserved in 70% solution of ethanol with a drop of glycerol added.

Diagnosis

Body length 630–645. Lamella absent. Rostral, lamellar, and interlamellar setae long, setiform, barbed; relative length *in* > *le* > *ro*. Bothridial seta short, clavate, barbed. Fourteen pairs of setae present: *c*₁ medium-sized, stiff, erect, barbed; *da*, *dm*, *dp*, *la*, *lm*, and *lp* long, setiform, barbed; *b*₁–*b*₃ and *p*₁–*p*₃ medium-sized, setiform, barbed. Notogastral porose area *Aa* elongate oval, others nearly circular. Epimeral and anogenital setae short, setiform, thin, nearly smooth or roughened. Leg chaetome complete.



Figures 1–3. *Phauloppia bageniaica* sp. nov. (adult) – 1. Dorsal view; 2. Ventral view (gnathosoma and legs omitted); 3. Right lateral view (gnathosoma and legs omitted).



Description

Integument – Body brown. Surface of body densely microgranulate sculpturing (clearer visible in humeral region).

Prodorsum – Rostrum rounded. Lamella completely absent, but several vaguely defined wrinkles present instead it. Rostral (82–90), lamellar (116–135), and interlamellar (139–157) setae setiform, barbed. Bothridial seta (36–45) clavate, barbed, with short stalk and head. Exobothridial seta (60–64) setiform, thin, slightly barbed. Dorsosejugal porose area oval (15–19 × 11–15).

Notogaster – Anterior margin distinctly convex medially. Fourteen pairs of setae present, c_1 (49–53) stiff, erect, barbed, others (da , dm , dp , la , lm , lp 101–116; c_2 , b_1 – b_3 75–86; p_1 – p_3 56–64) setiform, barbed. Four pairs of porose areas present, Aa elongate oval (30–34 × 11–13), $A1$, $A2$, and $A3$ nearly circular (11–19). Opisthonotal gland opening and all lyrifissures well visible.

Gnathosoma – Subcapitulum size 135–157 × 101–127. Three pairs of subcapitular setae (a 26–28; m 34–37; b 41–45) setiform, thin, roughened. Two pairs of adoral setae (15–17) setiform, thin, barbed. Palp length 97–101. Palp setal formula 0–2–1–3–9(+ ω). Postpalpal seta (7) spiniform, smooth. Chelicera length 143–161. Cheliceral setae (cha 45–49, chb 34) setiform, barbed.

Epimeral and lateral prodorsal regions – Epimeral setal formula 3–1–3–3. Epimeral setae $1a$, $2a$, and $3a$ (30–34) setiform, thin, nearly smooth; other setae (37–45) setiform, thin, roughened. Custodium absent. Discidium and circumpedal carina present. Humeral porose area Am and Ab oval.

Anogenital region – Anogenital setal formula 4–1–2–3. Genital (30–34), aggenital (37–45), anal (an_1 17–19; an_2 22–24), and adanal (ad_1 , ad_2 22–30; ad_3 30–34) setae setiform, thin, nearly smooth. Adanal lyrifissure oblique, located anterior to anal plate. Postanal porose area (visible in posterior view) band-like.

Legs – All leg tarsi tridactylous. Medial claw thicker than lateral claws. All claws slightly barbed on dorsal side. Lateral claws with small tooth distoventrally. Dorsal porose area on tarsus I, proximoventral porose area on tarsi I–IV, distoventral porose area on tibiae I–IV, and dorsoparaxial porose area on femora I–IV, and on trochanters III, IV distinct. Ventral porose area on tarsi and tibiae usually divided into two parts. Formulas of leg setation and solenidia: I (1–5–3–4–20) [1–2–2], II (1–5–3–4–15) [1–1–2], III (2–3–1–3–15) [1–1–0], IV (1–2–2–3–12) [0–1–0]. Homology of setae and solenidia indicated in Table 1.

Table 1. Leg setation and solenidia of adult *Phauloppia hageniaica* sp. nov.

Leg	Tr	Fe	Ge	Ti	Ta
I	v'	d , (l), bv'' , v''	(l), v' , σ	(l), (v), φ_1 , φ_2	(ft), (tc), (it), (p), (u), (a), s , (pv), v' , (pl), l'' , e , ω_1 , ω_2
II	v'	d , (l), bv'' , v''	(l), v' , σ	(l), (v), φ	(ft), (tc), (it), (p), (u), (a), s , (pv), ω_1 , ω_2
III	l' , v'	d , l' , ev'	l' , σ	l' , (v), φ	(ft), (tc), (it), (p), (u), (a), s , (pv)
IV	v'	d , ev'	d , l'	l' , (v), φ	ft'' , (tc), (p), (u), (a), s , (pv)

Note: Tr , Fe , Ge , Ti , Ta = trochanter, femur, genu, tibia, and tarsus, respectively. Roman letters refer to normal setae, Greek letters to solenidia, e = famulus. Single prime (') marks setae on the anterior and double prime (") setae on the posterior side of a given leg segment. Parentheses refer to a pair of setae.

Remarks

Phauloppia hageniaica sp. nov. is similar to *P. lucorum* (Koch, 1841) (supplementary descriptions see in, e.g., Mahunka 1982b; Weigmann 2006), from the Holarctic region and Nepal, in the length of the prodorsal and dorsal notogastral setae (relatively long) and the morphology of the notogastral seta c_1 (stiff and erect). However, the new species differs from the latter by the absence (versus presence) of the lamella and the morphology of the notogastral porose area Aa (elongate oval versus band-like).

Etymology

The specific name *hageniaica* refers to the tree (*Hagenia abyssinica*) on which the new species was found.

ACKNOWLEDGEMENTS

We thank Ethiopian colleagues, especially Dr. Gezahegn Degefe, for supporting field studies and

organizing laboratory work, and two anonymous reviewers for their valuable comments. The collection of materials was conducted under the Agreement between the Russian Academy of Sciences and the Ministry of Science and Technology (Ministry of Innovation and Technology) of the Federal Democratic Republic of Ethiopia.

Author contributions: All authors contributed equally on all counts. All authors have read and agreed to the published version of the manuscript.

Funding: This work was performed within the framework of the Joint Russian-Ethiopian Biological Expedition, financially supported by the Russian Academy of Sciences.

Data availability: Data are available upon request from the authors.

Ethics approval: This study only included arthropod material, and all required ethical guidelines for the treatment and use of animals were strictly adhered to in accordance with international, national, and institutional regulations. No human participants were involved in any studies conducted by the authors for this article.

Conflict of interest: The authors declare no conflict of interest.

Generative AI statement: The authors declare that AI was not used at any stage of the manuscript development.

REFERENCES

- Balogh, J. & Balogh, P. (2002) *Identification keys to the oribatid mites of the Extra-Holarctic regions*. Well-Press Publishing Limited, Miskolc, 453 pp. (Vol. 1) and 504 pp. (Vol. 2).
- Bayartogtokh, B. (2010) *Oribatid mites of Mongolia (Acari: Oribatida)*. KMK, Moscow, 372 pp.
- Berlese, A. (1895) *Acari, Myriapoda et Scorpiones hucusque in Italia reperta*. Portici, Padova, LXXVII, pp. 1–10.
- Berlese, A., (1896) *Acari, Myriapoda et Scorpiones hucusque in Italia reperta. Ordo Cryptostigmata II (Oribatidae)*. Portici, Padova, pp. 1–98.
- Berlese, A. (1908) Elenco di generi e specie nuove di Acari. *Redia*, 5: 1–15.
- Berlese, A. (1916) Centuria terza di Acari nuovi. *Redia*, 12: 289–338.
- Ermilov, S.G. & Rybalov, L.B. (2026) A checklist of the oribatid mite species (Acari, Oribatida) of Ethiopia. *Zootaxa*, 5773: 1–89. <https://doi.org/10.11646/zootaxa.5773.1.1>
- Ermilov, S.G. & Salavatulin, V.M. (2023) New arboreal mite species of Oribatulidae and Oripodidae (Acari, Oribatida, Oripodoidea) from Vietnam. *International Journal of Acarology*, 49: 283–291. <https://doi.org/10.1080/01647954.2023.2244979>
- Ermilov, S.G. & Salavatulin, V.M. (2025) New faunistic data on arboreal oribatid mites (Acari: Oribatida) of Vietnam. *Acarina*, 33: 3–7. <https://doi.org/10.21684/0132-8077-2025-33-1-3-7>
- Ermilov, S.G., Salavatulin, V.M. & Kontschán, J. (2024) Three new species of oribatid mites of the superfamily Oripodoidea (Acari, Oribatida) from Vietnam. *Zoologicheskij Zhurnal*, 103: 3–13. <https://doi.org/10.31857/S0044513424020011>
- Koch, C.L. (1841) *Deutschlands Crustaceen, Myriapoden und Arachniden, Heft 31*. F. Pustet, Regensburg.
- Mahunka, S. (1982a) Oribatids from the Eastern Part of the Ethiopian Region (Acari) I. *Acta Zoologica Academiae Scientiarum Hungaricae*, 28: 293–336.
- Mahunka, S. (1982b) Three new oribatid species (Acari) from Hungary. *Annales Historico-Naturales Musei Nationalis Hungarici*, 74: 295–299.
- Murvanidze, M. & Mumladze, L. (2016) Annotated checklist of Georgian oribatid mites. *Zootaxa*, 4089: 1–81. <https://doi.org/10.11646/zootaxa.4089.1.1>
- Norton, R.A. (1977) A review of F. Grandjean's system of leg chaetotaxy in the Oribatei (Acari) and its application to the family Damaeidae. In: Dindal, D.L., (Ed.), *Biology of oribatid mites*. SUNY College of Environmental Science and Forestry, Syracuse, pp. 33–61.
- Norton, R.A. & Behan-Pelletier, V.M. (2009) Suborder Oribatida. In: Krantz, G.W. & Walter, D.E. (Eds.),

- A manual of acarology*. Texas Tech University Press, Lubbock, USA, pp. 430–564.
- Salavatulin, V.M. (2019) Microhabitat distribution of arboreal oribatid mites (Oribatida), associated with the Siberian pine (*Pinus sibirica*) of Western Siberia. *Experimental and Applied Acarology*, 78: 469–483. <https://doi.org/10.1007/s10493-019-00401-4>
- Subías, L.S. (2022) Listado sistemático, sinonímico y biogeográfico de los ácaros oribátidos (Acariformes: Oribatida) del mundo (excepto fósiles). *Monografías Electrónicas Sociedad Entomológica Aragonesa*, 12: 1–538.
- Subías, L.S. (2024) Listado sistemático, sinonímico y biogeográfico de los ácaros oribátidos (Acariformes: Oribatida) del mundo (excepto fósiles), 19ª actualización: 1–545 (Accessed in January 2024).
- Thor, S. (1929) Über die Phylogenie und Systematik der Acarina, mit Beiträgen zur ersten Entwicklungsgeschichte einzelner Gruppen. *Nytt Magasin for Naturvidenskapene*, 67: 145–210.
- Travé, J. & Vachon, M. (1975) François Grandjean. 1882–1975 (Notice biographique & bibliographique). *Acarologia*, 17: 1–19.
- Weigmann, G. (2006) *Hornmilben (Oribatida)*. *Die Tierwelt Deutschlands*. Teil 76. Goecke and Evers, Keltern, 520 pp.

نخستین گزارش از جنس *Phauloppia* (Acari, Oribatida, Oribatulidae) در اتیوپی، به همراه توصیف گونه جدید درختی

سرگی جی. ارمیلوف^{۱*} | ولادیمیر ام. سالواتولین^۲ | لئونید بی. ریبالف^۲

۱. دانشگاه تیومن، مؤسسه زیست‌شناسی زیست‌محیطی و کشاورزی (X-BIO)، تیومن، روسیه؛ رایانامه: ermilovacari@yandex.ru

۲. مؤسسه اکولوژی و تکامل سورتسف، فرهنگستان علوم روسیه، آزمایشگاه جانورشناسی خاک و حشره‌شناسی عمومی، مسکو، روسیه؛ رایانامه‌ها: v.salavatulin@gmail.com و lybalov52@mail.ru

* نویسنده مسئول

✉ ermilovacari@yandex.ru

چکیده

جنس *Phauloppia* (Oribatida, Oribatulidae) برای نخستین بار از اتیوپی گزارش می‌شود. گونه جدیدی به نام *P. hageniaica* sp. nov. بر اساس کنه‌های ماده جمع‌آوری شده از پوست و خز درخت *Hagenia abyssinica* در جنوب شرقی اتیوپی توصیف می‌شود. این گونه جدید با نداشتن تیغه مشخص می‌شود.

واژگان کلیدی: ناحیه آفروتروپیکال، ریخت‌شناسی، هرناهای اربیاتید، آرایه‌شناسی، درخت‌زی

CITE: Ermilov, S.G., Salavatulin, V.M. & Rybalov, L.B. (2026) First record of the genus *Phauloppia* (Acari, Oribatida, Oribatulidae) in Ethiopia, with description of a new arboreal species. *Persian Journal of Acarology*, 15(3): 150310. <https://doi.org/10.22073/pja.150310>

دریافت

۴ اردیبهشت ۱۴۰۵

پذیرش

۱۹ خرداد ۱۴۰۵

انتشار

۲۴ تیر ۱۴۰۵

دبیر تخصصی

/. جوهرچی

