



Contribution to knowledge of the oribatid mite genus *Oxyamerus* (Acari, Oribatida, Oxyameridae), with description of a new species from Indonesia

Sergey G. Ermilov^{1*} | Dorothee Sandmann² | Stefan Scheu^{2,3}

1. University of Tyumen, X-BIO Institute, Tyumen, Russia; E-mail: ermilovacari@yandex.ru

2. University of Göttingen, JFB Institute of Zoology and Anthropology, Göttingen, Germany; E-mail: dsandma@gwdg.de, sscheu@gwdg.de

3. University of Göttingen, Centre of Biodiversity and Sustainable Land Use, Göttingen, Germany

* Correspondence

✉ ermilovacari@yandex.ru

Received:

27 July, 2026

Accepted:

25 August, 2026

Published:

15 October, 2026

Subject Editor:

O. Joharchi



<http://zoobank.org/urn:lsid:zoobank.org:pub:B743ADB6-D004-4E00-92DE-40F91D44843F>

ABSTRACT

The genus *Oxyamerus* (Oribatida, Oxyameridae) comprises 11 species distributed across the Australasian, Oriental, and southern Palearctic regions. A new species—*O. jambiensis* **sp. nov.**—is described, based on adults collected from forest litter in Sumatra, Indonesia. The new species is characterized by the presence of a long, thorn-like structure originating from the podosomal apophysis. An identification key, along with data on the distribution and habitats of the known representatives of *Oxyamerus*, are provided.

KEYWORDS

Generic diagnosis, identification key, Indonesia, morphology, oxyamerid mites, taxonomy

CITE: Ermilov, S.G., Sandmann, D. & Scheu, S. (2026) Contribution to knowledge of the oribatid mite genus *Oxyamerus* (Acari, Oribatida, Oxyameridae). *Persian Journal of Acarology*, 15(4): 150405.

<https://doi.org/10.22073/pja.150405>

INTRODUCTION

The oribatid mite genus *Oxyamerus* (Acari, Oribatida, Oxyameridae) was described by Aoki (1965), with *Oxyamerus spathulatus* Aoki, 1965 as type species. The genus currently contains 11 species which are collectively distributed in the Australasian, Oriental, and southern Palearctic regions (Subías 2022, online version 2024; Ermilov 2025). The generic diagnosis was revised by Ermilov and Corpuz-Raros (2017).

During the taxonomic identification of oribatid mites from Sumatra, Indonesia, one new species of *Oxyamerus* was found. The main goals of the paper are: (1) to describe this new species as *O. jambiensis* **sp. nov.**; (2) to provide an identification key for all known species of *Oxyamerus*; and (3) to compile data on the distribution and habitats of representatives of the genus.

Prior to this study, two species of *Oxyamerus*—*O. hyalinus* Hammer, 1979 and *O. truncatus* Hammer, 1979—were recorded from Indonesia (Hammer 1979).

MATERIAL AND METHODS

The specimens examined were collected in the framework of the EForTS Project ('Ecological and socioeconomic functions of tropical lowland rainforest transformation systems (Sumatra, Indonesia)'; <https://www.uni-goettingen.de/de/310995.html>; Drescher *et al.* 2016) on the island of Sumatra.

For measurement and illustration, specimens were mounted in lactic acid on temporary cavity slides. Body length was measured in lateral view, from the tip of the rostrum to the posterior edge of the



COPYRIGHT. Ermilov *et al.* Persian Journal of Acarology is under a free license. This open-access article is distributed under the terms of the [Creative Commons-BY-NC-ND](https://creativecommons.org/licenses/by-nc-nd/4.0/) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original author and source are credited.

gastronotum. Body width refers to the maximum width of the notogaster in dorsal view. Setae were measured from the side to avoid foreshortening. All measurements are given in micrometres (μm). Formulas for leg setation are given in parentheses according to the sequence trochanter–femur–genu–tibia–tarsus (famulus of tarsus I included). Formulas for leg solenidia are given in square brackets, according to the sequence genu–tibia–tarsus. Paired structures are described in the singular, unless otherwise noted. Drawings were made with a camera lucida using a Leica DM 2500 light microscope.

Morphological terminology used in this paper follows that of Grandjean: see Travé and Vachon (1975) for references, Norton (1977) for leg setal nomenclature, and Norton and Behan-Pelletier (2009) for overview.

Abbreviations: Prodorsum: *ro*, *le*, *in*, *bs*, *ex* = rostral, lamellar, interlamellar, bothridial, and exobothridial setae, respectively; *dst* = dorsosejugal tubercle. Notogaster: *c*, *la*, *lm*, *lp*, *b*₁–*b*₃, *p*₁–*p*₃ = setae; *ia*, *im*, *ip*, *ih*, *ips* = lyrifissures; *gla* = opisthonotal gland opening. Gnathosoma: *a*, *m*, *h* = subcapitular setae; *d*, *l*, *sup*, *inf*, *cm*, *acm*, *ul*, *su*, *lt*, *vt* = palp setae; ω = solenidion; *cha* = cheliceral seta; *Tg* = Trägårdh's organ. Epimeral and lateral podosomal regions: *1a*–*1c*, *2a*, *3a*–*3c*, *4a*–*4c* = epimeral setae; *PdI*, *PdII* = pedotecta I and II, respectively; *pap* = podosomal apophysis; *tls* = thorn-like structure. Anogenital region: *g*₁–*g*₆, *ag*, *an*₁–*an*₂, *ad*₁–*ad*₃ = genital, aggenital, anal, and adanal setae, respectively; *iad* = adanal lyrifissure; *po* = preanal organ. Legs: ω , φ , σ = solenidia; *e* = famulus; *d*, *l*, *v*, *bv*, *ev*, *ft*, *tc*, *it*, *p*, *u*, *a*, *s*, *pv*, *pl* = setae; *pa* = porose area.

RESULTS

TAXONOMY

Family Oxyameridae Aoki, 1965

Genus *Oxyamerus* Aoki, 1965

Type species: *Oxyamerus spatulatus* Aoki, 1965

Oxyamerus jambiensis sp. nov. (Figs 1–10)

<http://zoobank.org/urn:lsid:zoobank.org:act:AB8317DE-C0BD-45A2-8973-6A364F01CB3A>

Type material

Holotype (male) and two paratypes (one male and one female): Indonesia, Sumatra, Jambi Province, Bukit Duabelas National Park, 02° 05' 30.7" S, 102° 48' 30.7" E, litter in secondary rainforest, November 2013 (collected by B. Klarner).

Type deposition

The holotype is deposited in the collection of the Museum Zoologicum Bogoriense, Cibinong, Bogor, Indonesia; two paratypes are deposited in the collection of the University of Tyumen, Tyumen, Russia. All specimens are preserved in 70% ethanol with a drop of glycerol added.

Diagnosis


Body length 435–450. Body surface mostly densely microtuberculate. Rostrum rounded. Rostral seta setiform, indistinctly barbed. Lamellar seta setiform, thin, smooth. Interlamellar and exobothridial setae acicular. Relative length *ro* > *le* > *in* = *ex*. Bothridial seta long, nearly setiform, weakly expanded medially, with short cilia unilaterally. Dorsosejugal region with one pair of elongate tubercles. Humeral crista absent. All notogastral setae medium-sized, setiform, nearly smooth. Epimeral setal formula 3–1–3–3. Epimeral setae *1a*, *1b*, *1c*, *2a*, and *4a* acicular; other setae setiform, smooth; *3b* and *3c* longest. Podosomal apophysis large, nearly quadrangular. Strong thorn-like structure originating from podosomal apophysis developed. Three pairs of adanal setae.

Description

Sexual dimorphism – No significant sexual dimorphism was observed except body size.

Measurements – Body length 435 (male holotype), 435 (male paratype), 450 (female paratype).

Body width 213 (male holotype), 213 (male paratype), 225 (female paratype).




Figures 1–3. *Oxyamerus jambiensis* sp. nov. (adult) – 1. Dorsal view (legs except some trochanters omitted); 2. Ventral view (gnathosoma and legs except some trochanters omitted); 3. Right lateral view (gnathosoma and legs omitted).

Integument – Body color brown. Body covered by layer of gel-like cerotegument. Surface mostly densely microtuberculate, with weakly visible tubercles. Leg segments partially covered with simple or elongate distinct microtubercles.

Prodorsum – Rostrum rounded. Rostral seta (26–30) setiform, indistinctly barbed. Lamellar seta (13–15) setiform, thin, smooth. Interlamellar and exobothridial setae (7) acicular. Bothridial seta (82–86) nearly setiform, weakly expanded medially, with short cilia unilaterally. Dorsosejugal region (not humeral region) with one pair of strong, elongate, transversely oriented, simple tubercles located close and lateral

to bothridium.



Figures 4–10. *Oxyamerus jambiensis* sp. nov. (dissected adult) – 4. Subcapitulum, ventral view; 5. Palp, right, antiaxial view; 6. Chelicera, right, antiaxial view; 7. Leg I, right, antiaxial view; 8. Leg II, right, antiaxial view; 9. Leg III, left, turned; 10. Leg IV, right, turned.

Notogaster – Humeral crista absent. Ten pairs of notogastral setae (30–34) setiform, nearly smooth. Opisthonotal gland opening and all lyrifissures clearly visible.

Gnathosoma – Subcapitulum size 101–105 × 37. All subcapitular setae (*a*, *m* 15; *b* 11) setiform, indistinctly barbed. Palp length 71–75. Palp setal formula 0–2–1–3–9(+ω). Postpalpal seta absent. Chelicera length 101–109. Cheliceral seta *cha* (30–34) setiform, barbed; *chb* absent.

Epimeral and lateral podosomal regions – Epimeral setal formula 3–1–3–3. Setae *1a*, *1b*, *1c*, *2a*

(7–11), and *4a* (11–15) acicular. Other setae (*3b*, *3c* 71–75; *4c* 53–57; *4b* 41–45; *3a* 19–22) setiform, smooth. Setae *3b*, *4a*, and *4b* located close to each other. Podosomal apophysis large, nearly quadrangular, with weak lateral protrusion. Strong thorn-like structure originating from podosomal apophysis developed.

Anogenital region – Anogenital setal formula 6–1–2–3. Genital seta *g*₁ (19) setiform, thin, smooth; other genital setae (11), as well as aggenital (7), anal (7), and adanal (7) setae acicular. Adanal lyrifissure parallel and lateral to anal plate.

Legs – Claw of each leg strong, indistinctly barbed on dorsal side. Porose area on femora I–IV present, but poorly visible. Formulas of leg setation and solenidia: I (1–5–3–4–20) [1–2–2], II (1–5–3–4–16) [1–1–2], III (2–3–1–3–15) [1–1–0], IV (1–2–2–3–12) [0–1–0]. Homology of setae and solenidia indicated in Table 1. Setae *p'* and *p''* eupathidial (of tarsus I) or spiniform (tarsi II–IV). Seta *s* of tarsus I eupathidial. Some setae narrowly phylliform. Some solenidia slightly swollen distally.

Table 1. Leg setation of adult *Oxyamerus jambiensis* sp. nov.

Leg	Tr	Fe	Ge	Ti	Ta
I	<i>v'</i>	<i>d</i> , (<i>l</i>), <i>bv''</i> , <i>v''</i>	(<i>l</i>), <i>v'</i> , σ	(<i>l</i>), (<i>v</i>), φ_1 , φ_2	(<i>ft</i>), (<i>tc</i>), (<i>it</i>), (<i>p</i>), (<i>u</i>), (<i>a</i>), <i>s</i> , (<i>pv</i>), <i>v'</i> , (<i>pl</i>), <i>l''</i> , <i>e</i> , ω_1 , ω_2
II	<i>v'</i>	<i>d</i> , (<i>l</i>), <i>bv''</i> , <i>v''</i>	(<i>l</i>), <i>v'</i> , σ	(<i>l</i>), (<i>v</i>), φ	(<i>ft</i>), (<i>tc</i>), (<i>it</i>), (<i>p</i>), (<i>u</i>), (<i>a</i>), <i>s</i> , (<i>pv</i>), <i>l''</i> , ω_1 , ω_2
III	<i>l'</i> , <i>v'</i>	<i>d</i> , <i>l'</i> , <i>ev'</i>	<i>l'</i> , σ	<i>l'</i> , (<i>v</i>), φ	(<i>ft</i>), (<i>tc</i>), (<i>it</i>), (<i>p</i>), (<i>u</i>), (<i>a</i>), <i>s</i> , (<i>pv</i>)
IV	<i>v'</i>	<i>d</i> , <i>ev'</i>	<i>d</i> , <i>l'</i>	<i>l'</i> , (<i>v</i>), φ	<i>ft''</i> , (<i>tc</i>), (<i>p</i>), (<i>u</i>), (<i>a</i>), <i>s</i> , (<i>pv</i>)

Note: *Tr*, *Fe*, *Ge*, *Ti*, *Ta* = trochanter, femur, genu, tibia, and tarsus, respectively. Roman letters refer to normal setae, Greek letters to solenidia, *e* = famulus. Single prime (') marks setae on the anterior and double prime (") setae on the posterior side of a given leg segment. Parentheses refer to a pair of setae.

Remarks

Oxyamerus jambiensis sp. nov. differs from other species of the genus by the presence (vs. absence) of a strong thorn-like structure originating on the podosomal apophysis.

Etymology

The specific name *jambiensis* refers to the type locality, Jambi, Indonesia.

KEY TO KNOWN SPECIES OF *OXYAMERUS*

1. Rostral seta nearly setiform 2
- Rostral seta phylliform (dilated in middle or distal half) 3
2. All centrodorsal notogastral setae short (*la* about 1/2 of distance between *la* and *lp*). Notogastral seta *c* well developed, similar in length to centrodorsal notogastral setae. Podosomal apophysis with thorn-like structure. Second adanal seta inserted anterior to adanal lyrifissure *Oxyamerus jambiensis* sp. nov. Body length 435–450.
- All centrodorsal notogastral setae longer (*la* about 2/3 of distance between *la* and *lp*). Notogastral seta *c* not developed. Podosomal apophysis without thorn-like structure. Second adanal seta inserted lateral to adanal lyrifissure *Oxyamerus latirostris* Balogh, 1968. Body length 352.
3. Notogastral seta *c* well developed, similar in length to centrodorsal notogastral setae or slightly shorter 4
- Notogastral seta *c* short or not developed 6
4. All notogastral and second adanal setae bifurcate. Epimeral setae *1a*, *1b*, *1c*, *2a*, *3a*, *4a*, and aggenital setae phylliform. Lamellar seta short *Oxyamerus isabelaensis* Ermilov & Corpuz-Raros, 2017(a). Body length 448.
- All notogastral and second adanal setae not bifurcate. Epimeral setae *1a*, *1b*, *1c*, *2a*, *3a*, *4a*, and aggenital setae not phylliform. Lamellar seta medium-sized 5
5. Notogastral seta *c* bacilliform. Dorsosejugal tubercle (posterolateral to bothridium) absent. Epimeral setae *1a*, *1c*, *2a*, *3a*, and interlamellar seta spiniform. Adanal lyrifissure located lateral to anal aperture *Oxyamerus niedbalai* Ermilov & Kalúz, 2020. Body length 332–348.

- Notogastral seta *c* setiform. Dorsosejugal tubercle (posterolateral to bothridium) present. Epimeral setae *1a*, *1c*, *2a*, *3a*, and interlamellar seta setiform. Adanal lyrifissure located anterolateral to anal aperture *Oxyamerus konchuensis* Ermilov, 2025. Body length 360–390.
- 6. All centrodorsal notogastral setae short (*la* about 1/2 of distance between *la* and *lp*) 7
- All centrodorsal notogastral setae longer (*la* about 2/3 of distance between *la* and *lp* or longer)..... 9
- 7. Centrodorsal notogastral setae *la*, *lm*, *lp*, *b*₁–*b*₃ located on longitudinal granulate band. Bothridial seta smooth. Adanal lyrifissure located anterolateral to anal aperture
..... *Oxyamerus aoki* Balogh, 1968. Body length 279.
- Centrodorsal notogastral setae *la*, *lm*, *lp*, *b*₁–*b*₃ not located on longitudinal granulate band. Bothridial seta serrate unilaterally. Adanal lyrifissure located lateral to anal aperture 8
- 8. Rostral seta dilated medially, narrowed distally. Adanal lyrifissure oblique to anal aperture and distant from it *Oxyamerus himalayensis* Mondal & Kundu, 1999. Body length 430–442.
- Rostral seta dilated in distal half, rounded distally. Adanal lyrifissure close and parallel to anal aperture *Oxyamerus spathulatus* Aoki, 1965 (see also Aoki 1977). Body length 440–515.
- 9. Three pairs of adanal setae present
..... *Oxyamerus polilloensis* Corpuz-Raros & Lit, 2005. Body length 270–306.
- Two pairs of adanal setae present 10
- 10. First adanal seta inserted lateral to anal aperture. Notogastral seta *c* not developed. Notogastral setae *la* and *lm* located close to each other .. *Oxyamerus hauserorum* Mahunka, 1987. Body length 306–337.
- First adanal seta inserted posterior to anal aperture. Notogastral seta *c* well developed. Notogastral setae *la* and *lm* well separated from each other 11
- 11. Rostral seta broadly phylliform. Lamellar seta long
..... *Oxyamerus hyalinus* Hammer, 1979. Body length 360.
- Rostral seta narrowly phylliform. Lamellar seta medium-sized
..... *Oxyamerus truncatus* Hammer, 1979. Body length 470.

DISTRIBUTION AND HABITATS OF *OXYAMERUS*

Twelve species of *Oxyamerus* are distributed across the Australian, Oriental, and southern Palaearctic regions (Table 2). Six species have a highly circumscribed distribution, i.e. are endemic to a single country: *O. himalayensis* (India), *O. isabelaensis* (Philippines), *O. jambiensis* **sp. nov.** (Indonesia), *O. konchuensis* (Vietnam), *O. latirostris* (New Guinea), and *O. niedbalai* (Malaysia). Four species (*O. hauserorum*, *O. hyalinus*, *O. polilloensis*, and *O. truncatus*) are distributed in the Oriental region. *Oxyamerus aoki* is found in both Australian and Oriental regions; and *O. spathulatus* is registered in both Oriental and southern Palaearctic regions.

According to the summarized data (Table 2), *Oxyamerus* species predominantly inhabit forest litter and soil; however, some species have also been recorded in other microhabitats (e.g., moss, rotten wood).

Table 2. Distribution and habitats of *Oxyamerus* species.

Species	Indicated habitats	Countries	Main references
<i>O. aoki</i>	Ground moss. Litter in a mixed forest. Litter in forest concessions and commercial plantations. Soil in a mixed pine dominated forest. Soil in a mixed forest dominated by deciduous trees. Litter in forest	New Guinea, Philippines, southern China, Vietnam	Balogh 1968; Corpuz-Raros 1979; Corpuz-Raros & Lit 2013; Ermilov & Anichkin 2014; Ermilov 2025; Ni <i>et al.</i> 2025
<i>O. hauserorum</i>	Litter and soil in forest. Rotten wood in a dipterocarp forest	Borneo, Brunei, Malaysia	Mahunka 1987, 1988, 2005
<i>O. himalayensis</i>	Soil under <i>Camellia sinensis</i> . Soil under the decomposed leaves of <i>Dalbergia sissoo</i> . Soil under rotten leaves of <i>Shorea robusta</i> . Soil from loose humus. Soil with litter in forest	India	Mondal & Kundu 1999; Ermilov <i>et al.</i> 2014
<i>O. hyalinus</i>	Grass and semi-rotten hay on the roadside. The route follows an almost dry brook	Indonesia, Vietnam	Hammer 1979; Ermilov & Anichkin 2014

Table 2. Continued.

Species	Indicated habitats	Countries	Main references
	running through a luxurious forest with tree ferns, <i>Zingiber</i> sp., shrubs, dead trunks, mosses, dead leaves, and various debris. Soil in a mixed pine dominated forest. Soil in a mixed forest dominated by deciduous trees.		
<i>O. isabelaensis</i>	Litter under a large tree on a side rock of a cliff. Soil in the fork of a large tree, with little humus and ants. Bamboo litter	Philippines	Ermilov & Corpuz-Raros 2017a
<i>O. jambiensis</i>	Litter in forest	Indonesia	Present data
<i>O. konchuensis</i>	Litter in forest	Vietnam	Ermilov 2025
<i>O. latirostris</i>	Litter and decaying wood in jungle	New Guinea	Balogh 1968
<i>O. niedbalai</i>	Litter in forest	Malaysia	Ermilov & Kalúz 2020
<i>O. polilloensis</i>	Moss and decaying log. Decaying coconut trunk. Litter, soil, and moss in forest	Philippines, Sri Lanka	Corpuz-Raros & Lit 2005; Corpuz-Raros & Gruèzo 2005; Ermilov & Corpuz-Raros 2017b; Ermilov <i>et al.</i> 2019
<i>O. spathulatus</i>	Soil in forest	Thailand, Japan, Taiwan	Aoki 1965, 1977, 1998; Nakatamari 1983; Ermilov & Liao 2021
<i>O. truncatus</i>	The route follows an almost dry brook running through a luxurious forest with tree ferns, <i>Zingiber</i> sp., shrubs, dead trunks, mosses, dead leaves, and various debris. Soil in a mixed pine dominated forest. Soil in a mixed forest dominated by deciduous trees	Indonesia, Vietnam	Hammer 1979; Ermilov & Anichkin 2014

ACKNOWLEDGEMENTS

We thank two anonymous reviewers for their valuable comments; Dr. Rahayu Widyastuti (Institute Pertanian Bogor, Bogor, Indonesia) for her collaboration in this project; Dr. Bernhard Klarner (University of Göttingen, Göttingen, Germany) for collecting samples; Kristina Richter (University of Göttingen, Göttingen, Germany) for help in building up the Indonesian oribatid mite morphospecies collection and the database; the State Ministry of Research and Technology of Indonesia (RISTEK) for the research permit and the Indonesian Institute of Sciences (LIPI) and Ministry of Forestry (PHKA) for the collection permit; the Bukit Duabelas National Park for granting access to the field site; and the many colleagues and helpers for support in the field.

Author contributions: conceptualization, data curation, writing—original draft, visualization: SGE; investigation, writing—review and editing, methodology, data analysis, validation: SGE, DS, SS.

Funding: The study formed part of the German—Indonesian research project CRC990—EForTS funded by the German Research Foundation (DFG; project number 192626868-SFB 990).

Data availability: Data are available upon request from the author(s).

Ethics approval: This study only included arthropod material, and all required ethical guidelines for the treatment and use of animals were strictly adhered to in accordance with international, national, and institutional regulations. No human participants were involved in any studies conducted by the authors

for this article.

Conflict of interest: The authors declare no conflict of interest.

Generative AI statement: The authors declare that AI was not used at any stage of the manuscript development.

REFERENCES

- Aoki, J. (1965) Oribatiden (Acarina) Thailand. I. *Nature and Life in Southeast Asia*, 4: 129–193.
- Aoki, J. (1977) New and interesting species of oribatid mites from Kakeroma Island, Southwest Japan. *Acta Arachnologica*, 27: 85–93.
- Aoki, J. (1998) List of oribatid mites inhabiting forest soils of US Marine corps in Okinawa. *Bulletin Institute Environmental Science Technology*, 24: 141–145.
- Balogh, J. (1968) New oribatids (Acari) from New Guinea. *Acta Zoologica Academiae Scientiarum Hungaricae*, 14: 259–285.
- Corpuz-Raros, L.A. (1979) Philippine Oribatei (Acarina). I. Preliminary list of species and descriptions of forty new species. *Philippine Agriculturist*, 62: 1–82.
- Corpuz-Raros, L.A. & Gruezo, W.S. (2005) Preliminary list of soil mites (Acari) from the Samar Island Natural Park, Philippines with description of a new species and a new record of Oribatida. *Asia Life Sciences*, 14: 191–206.
- Corpuz-Raros, L.A. & Lit, I.L.jr. (2005) Some soil-inhabiting mites (Acari) from Polillo Island, Philippines, with descriptions of two new species of oribatids (Oribatida). *Asia Life Sciences*, 14: 179–190.
- Corpuz-Raros, L.A. & Lit, I.L.jr. (2013) Soil-inhabiting mites from the Quezon National Park and southern Sierra Madre Mountains in Quezon Province, Luzon Is., Philippines. *The Philippine Entomologist*, 27: 151–175.
- Drescher, J., Rembold, K., Allen, K., Beckschäfer, P., Buchori, D., Clough, Y., Faust, H., Fauzi, A.M., Gunawan, D., Hertel, D., Irawan, B., Jaya, I.N.S., Klarner, B., Kleinn, C., Knohl, A., Kotowska, M.M., Krashevskaya, V., Krishna, V., Leuschner, C., Lorenz, W., Meijide, A., Melati, D., Nomura, M., Pérez-Cruzado, C., Qaim, M., Siregar, I.Z., Steinebach, S., Tjoa, A., Tschardt, T., Wick, B., Wiegand, K., Kreft, H. & Scheu, S. (2016) Ecological and socioeconomic functions across tropical land-use systems after rainforest conversion. *Philosophical Transactions of the Royal Society B*, 371: 1–8. <https://doi.org/10.1098/rstb.2015.0275>
- Ermilov, S.G. (2025) Faunistic and taxonomic data on oribatid mites (Acari, Oribatida) from the Kon Chu Rang Nature Reserve, Vietnam. *Zoologicheskij Zhurnal*, 104: 3–14. <https://doi.org/10.7868/S3034543X25110017>
- Ermilov, S.G. & Anichkin, A.E. (2014) Taxonomic study of oribatid mites (Acari, Oribatida) of Bi Dup – Nui Ba National Park (southern Vietnam). *Zootaxa*, 3834: 1–86. <https://doi.org/10.11646/zootaxa.3834.1.1>
- Ermilov, S.G. & Corpuz-Raros, L. (2017a) New ameroid mites (Acari, Oribatida) from the Philippines. *Systematic and Applied Acarology*, 22: 1069–1086. <https://doi.org/10.11158/saa.22.7.14>
- Ermilov, S.G. & Corpuz-Raros, L. (2017b) New data on the Philippine oribatid mite fauna, with a contribution to knowledge of the genus *Drymobatoides* (Acari, Oribatida, Mochlozetidae). *Acarologia*, 57: 295–308. <https://doi.org/10.1051/acarologia/20174156>
- Ermilov, S.G. & Kalúz, S. (2020) New faunistic and taxonomic data on oribatid mites (Acari: Oribatida) of Malaysia. *Biologia*, 75: 1601–1611. <https://doi.org/10.2478/s11756-019-00411-y>
- Ermilov, S.G. & Liao, J.-R. (2021) New faunistic and taxonomic data on oribatid mites (Acari: Oribatida) of Taiwan. *Acarologia*, 61: 297–320. <https://doi.org/10.24349/acarologia/20214432>
- Ermilov, S.G., Kalúz, S., Martens, J. (2014) Additions to the Indian oribatid mite fauna, with description

- of a new species of the genus *Niphocephus* (Acari, Oribatida). *Systematic and Applied Acarology*, 19: 58–66. <https://doi.org/10.11158/saa.19.1.4>
- Ermilov, S.G., Khaustov, A.A. & Joharchi, O. (2019) New Oripodoidea (Acari, Oribatida) from Sri Lanka. *Systematic and Applied Acarology*, 24: 613–623. <https://doi.org/10.11158/saa.24.4.7>
- Hammer, M. (1979) Investigations on the oribatid fauna of Java. *Det Kongelige Danske Videnskaberne Selskab Biologiske Skrifter*, 22: 1–78.
- Mahunka, S. (1987) Neue und interessante Milben aus dem Genfer Museum LX. Oribatids from Sabah (East Malaysia) II. (Acari: Oribatida). *Revue Suisse de Zoologie*, 94: 765–817. <https://doi.org/10.5962/bhl.part.79551>
- Mahunka, S. (1988) New and interesting mites from the Geneva Museum LXI. Oribatids from Sabah (East Malaysia) III. (Acari: Oribatida). *Revue Suisse de Zoologie*, 95: 817–888. <https://doi.org/10.5962/bhl.part.81937>
- Mahunka, S. (2005) Oribatids from Brunei IV (Acari, Oribatida). (Acarologica Genavensia CVI). *Revue Suisse de Zoologie*, 112: 421–438. <https://doi.org/10.5962/bhl.part.80307>
- Mondal, B.K. & Kundu, B.G. (1999) A new oribatid mite (Acari) and four new records of oribatid from Indian soils. *Records of the Zoological Survey of India*, 97: 133–139.
- Nakatamari, S. (1983) The list of oribatid mites (Acari: Cryptostigmata) of the Ryukyu Islands, including a new record from Japan. *Biological Magazine Okinawa*, 21: 21–26.
- Ni, Z., Sun, J. & Liu, D. (2025) Oribatid mite (Acari, Oribatida) fauna of Bawangling National Nature Reserve in Hainan Island, China. *Systematic and Applied Acarology*, 30: 1846–1856. <https://doi.org/10.11158/saa.30.10.10>
- Norton, R.A. (1977) A review of F. Grandjean's system of leg chaetotaxy in the Oribatei (Acari) and its application to the family Damaeidae. In: Dindal, D.L. (Ed.), *Biology of oribatid mites*. SUNY College of Environmental Science and Forestry, Syracuse, pp. 33–61.
- Norton, R.A. & Behan-Pelletier, V.M. (2009) Suborder Oribatida. In: Krantz, G.W. & Walter, D.E. (Eds.), *A manual of acarology*. Texas Tech University Press, Lubbock, pp. 430–564.
- Subías, L.S. (2022) Listado sistemático, sinonímico y biogeográfico de los ácaros oribátidos (Acariformes: Oribatida) del mundo (excepto fósiles). *Monografías Electrónicas Sociedad Entomológica Aragonesa*, 12: 1–538.
- Subías, L.S. (2024) Listado sistemático, sinonímico y biogeográfico de los ácaros oribátidos (Acariformes: Oribatida) del mundo (excepto fósiles), 19ª actualización: 1–545 (Accessed on January 2024).
- Travé, J. & Vachon, M. (1975) François Grandjean. 1882–1975 (Notice biographique & bibliographique). *Acarologia*, 17: 1–19.

افزوده‌ای بر شناخت هرئای اورibatید جنس *Oxyamerus* (Acari, Oribatida, Oxyameridae) به همراه توصیف گونه جدیدی از اندونزی

سرگئی جی. ارمیلوف^{۱*} | دوروتی سندمن^۲ | استفان شو^۳

۱. دانشگاه تیومن، مؤسسه X-BIO، تیومن، روسیه؛ رایانامه: ermilovacari@yandex.ru
 ۲. دانشگاه گوتینگن، مؤسسه جانورشناسی و انسان‌شناسی JFB، گوتینگن، آلمان؛ رایانامه‌ها: sschen@gwdg.de dsandma@gwdg.de
 ۳. دانشگاه گوتینگن، مرکز تنوع زیستی و کاربری پایدار اراضی، گوتینگن، آلمان

* نویسنده مسئول

✉ usder@institutionaldomain

چکیده

جنس *Oxyamerus* (Oribatida, Oxyameridae) شامل ۱۱ گونه است که در مناطق استرالازیا، اورینتال (شرق آسیا) و بخش‌های جنوبی پالتارکتیک پراکندگی دارند. در این مقاله، گونه جدیدی با نام *O. jambiensis* sp. nov. بر اساس نمونه‌های کنه کامل جمع‌آوری شده از بستر جنگلی در سوماترا (اندونزی) توصیف می‌شود. ویژگی شاخص این گونه جدید، وجود ساختاری بلند و خارمانند است که از آپوفیز پاتنه‌ای (podosomal apophysis) منشأ می‌گیرد. همچنین، یک کلید شناسایی به همراه اطلاعاتی درباره پراکندگی و زیستگاه گونه‌های شناخته شده *Oxyamerus* ارائه شده است.

واژگان کلیدی: صفات مشخه جنس، کلید شناسایی، اندونزی، ریخت‌شناسی، کنه‌های اُکسیامرید، آرایه‌شناسی

CITE: Ermilov, S.G., Sandmann, D. & Scheu, S. (2026) Contribution to knowledge of the oribatid mite genus *Oxyamerus* (Acari, Oribatida, Oxyameridae). *Persian Journal of Acarology*, 15(4): 150405.
<https://doi.org/10.22073/pja.150405>

دریافت

۵ مرداد ۱۴۰۵

پذیرش

۳ شهریور ۱۴۰۵

انتشار

۲۳ مهر ۱۴۰۵

دبیر تخصصی

۱. جوهرچی

