



Persian J. Acarol., 2025, Vol. 14, No.3, pp. 465–470.
<https://doi.org/10.22073/pja.v14i3.87059>
Journal homepage: <http://www.biotaxa.org/pja>



Correspondence

The distribution and host selection of the chigger mite *Leptotrombidium yongshengense* (Acari: Trombiculidae) in southwest China

Chen-Xi Liu , Xian-Guo Guo , Yan Lv  and Peng-Wu Yin 

Institute of Pathogens and Vectors, Yunnan Provincial Key Laboratory for Zoonosis Control and Prevention, Dali University, Dali, Yunnan, 671000, China; E-mails: oOxcOo@163.com, xgguo2002@163.com, 867346563@qq.com, pengwuyin@vip.163.com

* Corresponding author

PAPER INFO.: Received: 23.03.2025, Accepted by: A. Saboori, 13.04.2025, Published: 15.07.2025

Chiggers (chigger mites) are the exclusive vector of *Orientia tsutsugamushi* (Ot), the causative agent of scrub typhus (tsutsugamushi disease). Although more than 3,000 chigger species have been recorded globally (Nielsen *et al.* 2021), only a few can effectively transmit Ot. Most effective vector species are from the genus *Leptotrombidium* of the family Trombiculidae (Paulraj *et al.* 2021; Balasubramanian *et al.* 2024), and *Leptotrombidium yongshengense* Yu and Yang, 1986 belongs to this genus. *Leptotrombidium yongshengense* was first named and described by Chinese scholars (Yu and Yang 1986), but little is known about this chigger species except the initial morphological description (Wang and Yu 1992; Stekolnikov 2013; Nielsen *et al.* 2021). Based on field surveys and taxonomic identification between 2001 and 2022, the present study retrospectively reports the distribution and host selection of *L. yongshengense* in the region for the first time.

The original data came from previous field surveys at 114 survey sites of southwest China (97° 21'–110 °11' E, 21° 08–33° 41' N) between 2001 and 2022. The 114 survey sites covered the five provincial regions of southwest China namely, Yunnan, Guizhou, Sichuan, Chongqing and Xizang (Tibet). Rodents and other sympatric small mammal hosts were routinely captured with mouse cages, and chiggers were collected from the body surface of the hosts (Li 1997; Guo *et al.* 2013). All the collected chiggers from each host were mounted onto glass slides with Hoyer's medium. After the process of drying, dehydration and transparency in an electric heat drying oven, all the chiggers were taxonomically identified to species under a microscope (Olympus, Tokyo, Japan) (Li 1997; Peng *et al.* 2015). Based on the taxonomic identification, *L. yongshengense* was selected as the subject of the present study. The constituent ratio (C_r), prevalence (P_M), mean abundance (MA) and mean intensity (MI) were calculated to reflect the infestation status of *L. yongshengense* on its hosts (Peng *et al.* 2018). The patchiness index (m^*/m) was used to measure the spatial distribution pattern of *L.*

How to cite: Liu, C.-X., Guo, X.-G., Lv, Y., Yin, P.-W. & Deak, G. (2025) The distribution and host selection of the chigger mite *Leptotrombidium yongshengense* (Acari: Trombiculidae) in southwest China. *Persian Journal of Acarology*, 14(3): 465–470.

In vertical distribution, *L. yongshengense* had relatively high C_r , P_M , MA and MI in high altitude gradients (1000–2000 m, 2000–3000 m, and > 3000 m) compared to the low altitude, under 1000 m ($P < 0.05$, Table 1). Most *L. yongshengense* came from the mountainous landscape ($C_r = 99.50\%$) with much higher infestation indices ($P_M = 2.59\%$, $MA = 0.49$ chiggers/per examined host, $MI = 18.90$ chiggers/per infested host) than those in the flatland landscape (Table 1, $P < 0.01$). All 7,028 *L. yongshengense* were from the outdoor habitat, and no *L. yongshengense* was found in the indoor habitat. The results reveal the fluctuation of *L. yongshengense* in different environments with environmental heterogeneity. *Leptotrombidium yongshengense* have a tendency to live in outdoors of mountainous areas with high altitudes.

Table 1. Geographical distribution of *Leptotrombidium yongshengense* along different environment gradients in southwest China (2001–2022).

Different environmental gradients		<i>L. yongshengense</i>		Infestation indices		
		No.	C_r (%)	P_M (%)	MA	MI
Altitudes (m)	< 1000	42	0.6	0.35	0.01	2.47
	1000–2000	3579	50.92	2.3	0.56	24.35
	2000–3000	2508	35.69	2.68	0.38	14.17
	> 3000	899	12.79	1.23	0.29	23.66
Landscape	Flatland	35	0.5	0.14	0.01	3.89
	Mountainous	6993	99.5	2.59	0.49	18.9

The majority of *L. yongshengense* were identified from *Apodemus chevrieri* ($C_r = 36.95\%$, 2597/7028), *Tupaia belangeri* ($C_r = 31.76\%$, 2232/7028) and *Alexandromys oeconomus* ($C_r = 11.23\%$, 789/7028), with the highest infestation indices on the vole *A. oeconomus* (Table 2, $P < 0.05$). The results suggest that *L. yongshengense* has an obvious host-preference, and *A. oeconomus* has a high susceptibility to the infestation of *L. yongshengense*. A previous report showed that *L. deliense*, belonging to the same genus with *L. yongshengense*, was mainly distributed in low altitude areas in southwest China, and its main host was the rat *Rattus tanezumi* (Lv et al. 2021). The above differences between *L. yongshengense* and *L. deliense* reflect the interspecific difference in geographical distribution and host selection.

Table 2. Infestation indices and patchiness index (m^*/m) of *Leptotrombidium yongshengense* on three important species of small mammal hosts in southwest China (2001–2022).

Names of hosts	Constituent ratios and infestation indices of <i>L. yongshengense</i>					m^*/m
	No.	C_r , %	P_M , %	MA	MI	
<i>A. chevrieri</i>	2597	36.95	4.14	1.09	26.23	168.26
<i>T. belangeri</i>	2232	31.76	28.88	8.06	27.90	11.13
<i>A. oeconomus</i>	789	11.23	58.33	65.75	112.71	2.89

The values of m^*/m for *L. yongshengense* were higher than the border value ($m^*/m > 1$) for determining the aggregated distribution (Table 2), indicating *L. yongshengense* is of aggregated distribution among different individuals of its hosts. The aggregated distribution pattern of *L.*

yongshengense is consistent with other chigger species (Yin *et al.* 2021), which is beneficial to the growth, development, reproduction and population expansion of chiggers and some other parasites (Nicácio *et al.* 2019). The association coefficient for *L. yongshengense* and *L. rusticum* was $V = 0.10$ ($\chi^2 = 216.47$, $P < 0.01$), and for *L. yongshengense* and *L. scutellare* was $V = 0.07$ ($\chi^2 = 104.79$, $P < 0.01$). Two association coefficients were very low (close to 0), indicating that these chigger species are almost independent of each other in host selection (Guo *et al.* 2006).

Most *L. yongshengense* were from adult hosts ($C_r = 97.20\%$, 6831/7028) with higher P_M , MA , and MI than those on juvenile hosts (Table 3, $P < 0.05$), indicating that *L. yongshengense* tends to parasitize adult hosts with age-bias in the selection of hosts. The large body surface area, frequent foraging and activities may make adult hosts easy to be invaded by chigger mites and other parasites (Kataranovski *et al.* 2011). More of *L. yongshengense* were found on male hosts; however, the differences of P_M , MA and MI between male and female hosts were of no statistical significance (Table 3, $P > 0.05$).

Table 3. Infestation indices of *Leptotrombidium yongshengense* on different ages and sexes of hosts in southwestern China (2001–2022).

Ages and sexes of hosts		Constituent ratios and infestation indices of <i>L. yongshengense</i>				
		No.	C_r (%)	P_M (%)	MA	MI
Ages	Adult	6831	97.20	2.51	0.49	19.35
	Juvenile	197	2.80	0.81	0.06	7.58
Sexes	Female	3343	47.57	2.11	0.41	19.44
	Male	3684	52.43	2.27	0.41	17.88

The present study indicates that *L. yongshengense* is a common species of chigger mites with a wide geographical distribution and large population in southwest China. The study has enriched the scientific information about *L. yongshengense* and the genus *Leptotrombidium*. Due to the lack of direct evidence to prove whether *L. yongshengense* can effectively transmit Ot, the epidemiological significance of its distribution in southwest China remains to be further studied.

REFERENCES

Balasubramanian, T., Sambath, U., Radja, R.D., Thangaraj, G., Devaraju, P., Srinivasan, L., Srinivasa, P., Nair, M.G., Raja, K., Lakkawar, A.W. & Soong, L. (2024) Pathological responses in Asian house shrews (*Suncus murinus*) to the naturally acquired *Orientia tsutsugamushi* infection. *Microorganisms*, 12(4): 748. DOI: [10.3390/microorganisms12040748](https://doi.org/10.3390/microorganisms12040748)

Guo, X.G., Qian, T.J., Meng, X.Y., Dong, W.G., Wu, D. & Shi, W.X. (2006) Preliminary analysis of chigger communities associated with house rats (*Rattus flavipectus*) from six counties in Yunnan, China. *Systematic & Applied Acarology*, 11(1): 13–21. DOI: [10.11158/saa.11.1.2](https://doi.org/10.11158/saa.11.1.2)

Guo, X.G., Speakman, J.R., Dong, W.G., Men, X.Y., Qian, T.J., Wu, D., Qin, F. & Song, W.Y. (2013) Ectoparasitic insects and mites on Yunnan red-backed voles (*Eothenomys miletus*) from a

- localized area in southwest China. *Parasitology Research*, 112(10): 3543–3549. DOI: [10.1007/s00436-013-3537-6](https://doi.org/10.1007/s00436-013-3537-6)
- Kataranovski, M., Mirkov, I., Belij, S., Popov, A., Petrović, Z., Gačić, Z. & Kataranovski, D. (2011) Intestinal helminths infection of rats (*Ratus norvegicus*) in the Belgrade area (Serbia): The effect of sex, age and habitat. *Parasite*, 18(2): 189–196. DOI: [10.1051/parasite/2011182189](https://doi.org/10.1051/parasite/2011182189)
- Li, J.C. (1997) *Trombiculid mites of China: Studies on vector and pathogen of tsutsugamushi disease*. First edition. Guangdong Science and Technology Press, Guangzhou, China, 570 pp. (in Chinese)
- Lopez, J.E. (2005) Parasite prevalence and the size of host populations: An experimental test. *Journal of Parasitology*, 91(1): 32–37. DOI: [10.1645/GE-323R](https://doi.org/10.1645/GE-323R)
- Lv, Y., Guo, X.G., Jin, D.C., Fan, R., Zhao, C.F., Zhang, Z.W., Mao, K.Y., Song, W.Y., Dong, W.G., Qian, T.J. & Yang, Z.H. (2021) Distribution and host selection of the chigger mite vector of scrub typhus, *Leptotrombidium deliense*, in southwest China. *International Journal of Acarology*, 47(3): 233–241. DOI: [10.1080/01647954.2021.1890210](https://doi.org/10.1080/01647954.2021.1890210)
- Nicácio, J., Oliveira, I.D., Uchoa, M.A., Faccenda, O., Abot, A.R., Fernandes, M.G. & Garcia, F.R. (2019) Spatial distribution and control levels of *Anastrepha* spp. (Diptera: Tephritidae) in guava orchards. *Annals of the Brazilian Academy of Sciences*, 91(03): e20180428. DOI: [10.1590/0001-3765201920180428](https://doi.org/10.1590/0001-3765201920180428)
- Nielsen, D.H., Robbins, R.G. & Rueda, L.M. (2021) Annotated world checklist of the Trombiculidae and Leeuwenhoeekiidae (1758–2021) (Acari: Trombiculoidea), with notes on nomenclature, taxonomy, and distribution. *Zootaxa*, 4967(1): 001–243. DOI: [10.11646/zootaxa.4967.1.1](https://doi.org/10.11646/zootaxa.4967.1.1)
- Paulraj, P.S., Renu, G., Ranganathan, K., Leo, V.J. & Veeramanoharan, R. (2021) First seroprevalence report of scrub typhus from the tribal belts of the Nilgiris district, Tamil Nadu, India. *Indian Journal of Medical Research*, 153(4): 503–507. DOI: [10.4103/ijmr.IJMR_1223_19](https://doi.org/10.4103/ijmr.IJMR_1223_19)
- Peng, P.Y., Guo, X.G., Song, W.Y., Hou, P., Zou, Y.J., Fan, R. & He, X.S. (2015) Analysis of ectoparasites (chigger mites, gamasid mites, fleas and sucking lice) of the Yunnan red-backed vole (*Eothenomys miletus*) sampled throughout its range in southwest China. *Medical and Veterinary Entomology*, 29(4): 403–415. DOI: [10.1111/mve.12134](https://doi.org/10.1111/mve.12134)
- Peng, P.Y., Guo, X.G., Jin, D.C., Dong, W.G., Qian, T.J., Qin, F., Yang, Z.H. & Fan, R. (2018) Landscapes with different biodiversity influence distribution of small mammals and their ectoparasitic chigger mites: A comparative study from southwest China. *PLoS ONE*, 13(1): e0189987. DOI: [10.1371/journal.pone.0189987](https://doi.org/10.1371/journal.pone.0189987)
- Stekolnikov, A.A. (2013) *Leptotrombidium* (Acari: Trombiculidae) of the World. *Zootaxa*, 3728(1): 001–173. DOI: [10.11646/zootaxa.3728.1.1](https://doi.org/10.11646/zootaxa.3728.1.1)
- Wang, D.Q. & Yu, Z.Z. (1992) Chigger mites of the genus *Leptotrombidium*: Key to species and their distribution in China. *Medical and Veterinary Entomology*, 6(4): 389–395. DOI: [10.1111/j.1365-2915.1992.tb00639.x](https://doi.org/10.1111/j.1365-2915.1992.tb00639.x)
- Yin, P.W., Guo, X.G., Jin, D.C., Fan, R., Zhao, C.F., Zhang, Z.W., Huang, X.B. & Mao, K.Y. (2021) Distribution and host selection of tropical rat mite, *Ornithonyssus bacoti*, in Yunnan Province of southwest China. *Animals*, 11(1): 110. DOI: [10.3390/ani11010110](https://doi.org/10.3390/ani11010110)

Yu, Z.Z. & Yang, G.R. (1986) Five new species of chigger mites from western part of Yunnan Province (Acariformes: Trombiculidae). *Acta Zootaxonomica Sinica*, 11: 63–71.

COPYRIGHT

Liu *et al.* Persian Journal of Acarology is under a free license. This open-access article is distributed under the terms of the Creative Commons-BY-NC-ND which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original author and source are credited.